

## CONCLUSION

The incidental finding of focal thyroid uptake on FDG-PET scanning, a so-called PEToma, warrants careful evaluation as a substantial number will harbour a primary thyroid cancer. Use of maximum SUV values alone is a poor predictor of malignancy in these patients. At a minimum, these patients require further evaluation with ultrasonography and FNAC.

**Acknowledgements:** We thank Richard J. Payne, MD, of the Department of Otolaryngology, Royal Victoria Hospital, Montréal, for his scientific advice and Shinichi Hisasue, MD, of the Department of Psychology, University of Massachusetts, for his assistance in the statistical analysis.

**Competing interests:** None declared.

**Contributors:** Drs. Tabah and How designed the study. Drs. Nishimori and How wrote the article, which Drs. Tabah, Hickeson and How critically reviewed. All authors acquired and analyzed the data and approved publication of the article.

## References

- Davies L, Welch HG. Increasing incidence of thyroid cancer in the United States, 1973-2002. *JAMA* 2006;295:2164-7.
- How J, Tabah R. Explaining the increasing incidence of differentiated thyroid cancer. *CMAJ* 2007;177:1383-4.
- Are C, Hsu JF, Schoder H, et al. FDG-PET detected thyroid incidentalomas: Need for further investigation? *Ann Surg Oncol* 2007;14:239-47.
- Salvatori M, Melis L, Castaldi P, et al. Clinical significance of focal and diffuse thyroid disease identified by <sup>18</sup>F-fluorodeoxyglucose positron emission tomography. *Biomed Pharmacother* 2007;61:477-81.
- Are C, Hsu JF, Chossein RA, et al. Histological aggressiveness of fluorodeoxyglucose positron-emission tomogram (FDG-PET)-detected incidental thyroid carcinomas. *Ann Surg Oncol* 2007;14:3210-5.
- Cohen MS, Arslan N, Dehdashti F, et al. Risk of malignancy in thyroid incidentalomas identified by fluorodeoxyglucose-positron emission tomography. *Surgery* 2001;130:941-6.
- Chen YK, Ding HJ, Chen KT, et al. Prevalence and risk of cancer of focal thyroid incidentaloma identified by <sup>18</sup>F-fluorodeoxyglucose positron emission tomography for cancer screening in healthy subjects. *Anticancer Res* 2005;25:1421-6.
- Choi JY, Lee KS, Kim HJ, et al. Focal thyroid lesions incidentally identified by integrated <sup>18</sup>F-FDG PET/CT: clinical significance and improved characterization. *J Nucl Med* 2006;47:609-15.
- Kim TY, Kim WB, Ryu JS, et al. <sup>18</sup>F-fluorodeoxyglucose uptake in thyroid from positron emission tomogram (PET) for evaluation in cancer patients: high prevalence of malignancy in thyroid PET incidentaloma. *Laryngoscope* 2005;115:1074-8.
- Bogsrud TV, Karantanis D, Nathan MA, et al. The value of quantifying <sup>18</sup>F-FDG uptake in thyroid nodules found incidentally on whole-body PET-CT. *Nucl Med Commun* 2007;28:373-81.
- Chu QD, Connor MS, Lilien DL, et al. Positron emission tomography (PET) positive thyroid incidentaloma: the risk of malignancy observed in a tertiary referral center. *Am Surg* 2006;72:272-5.
- Cooper DS, Doherty GM, Haugen BR, et al. Management guidelines for patients with thyroid nodules and differentiated thyroid cancer. *Thyroid* 2006;16:109-42.
- NCCN Clinical Practice Guidelines in Oncology. In: National Comprehensive Cancer Network website. Available: [www.nccn.org/professionals/physician\\_gls/f\\_guidelines.asp](http://www.nccn.org/professionals/physician_gls/f_guidelines.asp) (accessed 2010 Dec. 8).
- Kang HW, No JH, Chung JH, et al. Prevalence, clinical and ultrasonographic characteristics of thyroid incidentalomas. *Thyroid* 2004;14:29-33.
- Burguera B, Gharib H. Thyroid incidentalomas. Prevalence, diagnosis, significance, and management. *Endocrinol Metab Clin North Am* 2000;29:187-203.
- King DL, Stack BC Jr, Spring PM, et al. Incidence of thyroid carcinoma in fluorodeoxyglucose positron emission tomography-positive thyroid incidentalomas. *Otolaryngol Head Neck Surg* 2007;137:400-4.
- Shie P, Cardarelli R, Sprawls K, et al. Systematic review: prevalence of malignant incidental thyroid nodules identified on fluorine-18 fluorodeoxyglucose positron emission tomography. *Nucl Med Commun* 2009;30:742-8.
- Mitchell JC, Grant F, Evenson AR, et al. Preoperative evaluation of thyroid nodules with <sup>18</sup>FDG-PET/CT. *Surgery* 2005;138:1166-74.
- Rosenbaum SJ, Lind T, Antoch G, et al. False-positive FDG PET uptake — the role of PET/CT. *Eur Radiol* 2006;16:1054-65.

## CORRECTION

In the February issue of the *Canadian Journal of Surgery*, incorrect data were reported in Table 1 in the following research paper: Zahrai A, Chahal J, Stojimirovic D, et al. Quality of life and educational benefit among orthopedic surgery residents: a prospective, multicentre comparison of the night float and the standard call systems. *Can J Surg* 2011;54(1):25-32. This error was in the print version of the journal only; the online version has been corrected.