

- cancer: psychological morbidity. *Eur J Cancer* 1992;28A(8-9):1340-4.
43. Schain WS, d'Angelo TM, Dunn ME, Lichter AS, Pierce LJ. Mastectomy versus conservative surgery and radiation therapy. Psychosocial consequences. *Cancer* 1994; 73:1221-8.
 44. de Haes JC, Curran D, Aaronson NK, Fentiman IS. Quality of life in breast cancer patients aged over 70 years, participating in the EORTC 10850 randomised clinical trial. *Eur J Cancer* 2003;39:945-51.
 45. Moyer A. Psychosocial outcomes of breast-conserving surgery versus mastectomy: a meta-analytic review. *Health Psychol* 1997;16:284-98.
 46. Fisher B, Jeong JH, Anderson S, Bryant J, Fisher ER, Wolmark N. Twenty-five-year follow-up of a randomized trial comparing radical mastectomy, total mastectomy, and total mastectomy followed by irradiation. *N Engl J Med* 2002;347:567-75.
 47. Louis-Sylvestre C, Clough K, Asselain B, Vilcoq JR, Salmon RJ, Campana F, et al. Axillary treatment in conservative management of operable breast cancer: Dissection or radiotherapy? Results of a randomized study with 15 years of follow-up. *J Clin Oncol* 2004;22:97-101.
 48. Phase III randomized study of sentinel node dissection with or without conventional axillary dissection in women with clinically node negative breast cancer. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrd=66987&protocolsearchid=917856 (accessed 2005 Mar 30).
 49. Phase III prognostic study of sentinel node and bone marrow micrometastases in women with stage I or IIA breast cancer. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrd=67017&protocolsearchid=917814 (accessed 2005 Mar 30).
 50. Phase III randomized study of axillary lymph node dissection in women with stage I or IIA breast cancer who have a positive sentinel node. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrd=67018&protocolsearchid=917762 (accessed 2005 Mar 30).
 51. Phase III randomized study of surgical resection with or without axillary lymph node dissection in women with clinically node-negative breast cancer with sentinel node micrometastases. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrd=339581&protocolsearchid=917850 (accessed 2005 Mar 30).
 52. Phase III randomized study of complete axillary lymph node dissection versus axillary radiotherapy in sentinel lymph node-positive women with operable invasive breast cancer. CANCERNET-PDQ [database online]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrd=68566&protocolsearchid=917889 (accessed 2005 Mar 30).
 53. Cantin J, Scarth H, Levine M, Hugi M. Clinical practice guidelines for the care and treatment of breast cancer: 13. Sentinel lymph node biopsy. *CMAJ* 2001; 165:166-73.
 54. Powles TJ, Hickish TF, Makris A, Ashley SE, O'Brien ME, Tidy VA, et al. Randomized trial of chemoendocrine therapy started before or after surgery for treatment of primary breast cancer. *J Clin Oncol* 1995;13:547-52.
 55. Wolmark N, Wang J, Mamounas E, Bryant J, Fisher B. Preoperative chemotherapy in patients with operable breast cancer: nine-year results from National Surgical Adjuvant Breast and Bowel Project B-18. *J Natl Cancer Inst Monogr* 2001;(30):96-102.
 56. van der Hage JA, van de Velde CJ, Julien JP, Tubiana-Hulin M, Vandervelden C, Duchateau L. Preoperative chemotherapy in primary operable breast cancer: results from the European Organization for Research and Treatment of Cancer trial 10902. *J Clin Oncol* 2001;19:4224-37.
 57. Gianni L, Baselga J, Eirmann W, Porta VG, Semiglazov V, Garcia-Conde J, et al. First report of the European Cooperative Trial in operable breast cancer (ECTO): effects of primary systemic therapy (PST) on local-regional disease [abstract]. *Proc Am Soc Clin Oncol* 2002;21(1):34a.
 58. Phase III randomized study of surgery with or without axillary node clearance followed by adjuvant tamoxifen in elderly women with breast cancer. CANCERNET-PDQ [online database]. Available: www.cancer.gov/search/ViewClinicalTrials.aspx?cdid=78383&version=HealthProfessional&protocolsearchid=1008021 (accessed 2005 Mar 30).
 59. Harris SR, Hugi MR, Olivotto IA, Levine M. Clinical practice guidelines for the care and treatment of breast cancer: 11. Lymphedema. *CMAJ* 2001;164:191-9.
 60. Tengrup I, Tennvall-Nittby L, Christiansson I, Laurin M. Arm morbidity after breast-conserving therapy for breast cancer. *Acta Oncol* 2000;39:393-7.
 61. McCready DR, Bodurtha AJ, Davis NL, Meterissian S, Robert J, Temple WJ. Sentinel lymph-node biopsy in breast cancer. *Can J Surg* 1999;42:406-7.

Correction

In a case report in the April 2005 issue of the journal (*Can J Surg* 2005;48[2]:159-60), one of the authors' names was misspelled. The author byline on page 159 should read as follows: Shahzeer Karmali, MD, BSc; * Luke Rudmik, MD; * Walley Temple, MD; * Vincent Falck, MB ChB; † Gregory McKinnon, MD*