

Prospective clinical and radiographic results of CHARITÉ III artificial total disc arthroplasty at 2- to 7-year follow-up: a Canadian experience

Michael Katsimihas, MSc,
MBChB

Christopher S. Bailey, MD, MSc
Khalil Issa, MD

Jennifer Fleming, PhD

Patricia Rosas-Arellano, MD,
MSc, PhD

Stewart I. Bailey, MD

Kevin R. Gurr, MD

From the Division of Orthopaedics,
Department of Surgery, University of
Western Ontario, and the Orthopaedic
Spine Program, Victoria Hospital, Lon-
don Health Science Centre, London, Ont.

This work was presented at the meeting
of the British Association of Spine
Surgeons, Sheffield, England, Feb. 25–27,
2009, and at the Annual Meeting of the
Canadian Orthopaedic Association,
Whistler, BC, July 3–6, 2009.

Accepted for publication
Mar. 31, 2010

Correspondence to:

Dr. C.S. Bailey
London Health Science Centre
Victoria Hospital
800 Commissioners Rd. E, E4 120
London ON N6A 5W9
chris.bailey@lhsc.on.ca

Background: Early and intermediate results have shown that the SB CHARITÉ III total disc arthroplasty (TDA) favourably compares to spinal fusion, but is associated with fewer complications and higher levels of satisfaction. We sought to prospectively report the clinical and radiographic results of the CHARITÉ III TDA after an average of 55 months follow-up.

Methods: We conducted a prospective study of patients receiving the CHARITÉ TDA at either L4–5 or L5–S1 between April 2001 and November 2006. The primary indication for surgery was discogenic low-back pain confirmed by provocative discography. Assessment included pre- and postoperative (3, 6 and 12 mo and yearly thereafter) validated patient outcome measures and radiographic review.

Results: Fifty-seven of the potential 64 (89%) patients were available for complete follow-up. Their mean age was 39 (range 21–59) years. A statistically significant improvement was demonstrated between all the mean pre- and postoperative intervals for the Oswestry Disability Index, visual analogue scale for back and leg pain, and Short Form-36 health survey ($p < 0.001$). The mean sagittal rotation was 6.5° (range 0.5° – 22.4°), and the mean intervertebral translation was 1.1 mm (range 0–2.4 mm). Subsidence of the implant was present in 44 of 53 (83%) patients with an L5–S1 disc arthroplasty. The mean subsidence was 1.7 mm (range 0–4.8 mm).

Conclusion: The 2- to 7-year follow-up of this cohort of patients demonstrated satisfactory clinical and radiographic results in a carefully selected patient population. The radiographic assessment confirmed preservation and maintenance of motion at the replaced disc during the period of follow-up.

Contexte : Les premiers résultats et les résultats intermédiaires ont démontré que l'arthroplastie discale totale (ADT) SB CHARITÉ III se compare favorablement à l'arthrodèse de la colonne, mais qu'on y associe moins de complications et une plus grande satisfaction. Nous avons cherché à faire état de façon prospective des résultats cliniques et radiographiques de l'ADT CHARITÉ III après un suivi d'une durée moyenne de 55 mois.

Méthodes : Nous avons effectué une étude prospective portant sur des patients qui ont subi une ADT CHARITÉ au niveau L4–5 ou L5–S1 entre avril 2001 et novembre 2006. L'indication principale pour l'intervention chirurgicale était une lombalgie d'origine discale confirmée par discographie de provocation. L'évaluation a inclus des mesures de résultats validées avant et après l'intervention (3, 6 et 12 mois et une fois l'an par la suite) et un examen radiographique.

Résultats : Sur les 64 patients possibles, 59 (92 %) étaient disponibles pour un suivi complet. Ils avaient en moyenne 39 ans (intervalle de 21–59 ans). On a démontré une amélioration statistiquement significative entre tous les intervalles moyens avant et après l'intervention pour ce qui est de l'évaluation de l'incapacité fonctionnelle Oswestry, de l'échelle analogique visuelle pour la douleur au dos et à la jambe, et du questionnaire généraliste SF-36 (Short Form-36) sur la santé ($p < 0,001$). La rotation sagittale moyenne s'établissait à $6,5^\circ$ (intervalle de $0,5^\circ$ – $22,4^\circ$) et la translation intervertébrale moyenne, à 1,1 mm (intervalle de 0–2,4 mm). Il y avait régression de l'implant chez 44 patients (83 %) sur 53 qui avaient subi une arthroplastie discale L5–S1. La régression moyenne s'établissait à 1,7 mm (intervalle de 0–4,8 mm).

Conclusion : Le suivi d'une durée de 2 à 7 ans de cette cohorte de patients a démontré des résultats cliniques et radiographiques satisfaisants dans une population de patients sélectionnés avec soin. L'évaluation radiographique a confirmé la conservation et le maintien du mouvement au niveau du disque remplacé au cours de la période de suivi.

Lumbar spine fusion remains a treatment option for the management of patients with low-back pain due to degenerative disc disease for whom nonoperative options have failed. However, spinal fusion has been related to the development of facet joint arthrosis, degeneration of the discs adjacent to the fusion, spinal stenosis, instability and dysfunction of the sacroiliac joints.⁷⁻¹¹ In a study of 215 patients who underwent lumbar fusion, a reoperation rate of 27% was reported at a mean follow-up of 6.7 years for adjacent level disease.⁸ Similarly, in another study, adjacent level disease resulted in a 20% reoperation rate after 2–15 years of follow-up.⁹

To prevent possible consequences of lumbar fusion, the total disc arthroplasty (TDA) was introduced as an alternative to fusion. It is theorized that an artificial disc replacement preserves motion at the operated levels and reduces the strain on adjacent levels, thus minimizing the chance for development of adjacent level disease. This theory has been supported in biomechanical cadaveric studies; the CHARITÉ TDA was shown to restore motion to the level of the intact segment in flexion–extension, lateral flexion and axial rotation, reproducing a motion very similar to the physiologic motion of an intervertebral disc.^{13,14} A number of clinical studies also demonstrated preservation of motion at the operated level.^{15,16}

Early and intermediate results have shown that the SB CHARITÉ III TDA (DePuy Spine) favourably compares with spinal fusion and is associated with fewer complications and higher levels of satisfaction than fusion.¹⁶⁻²² Limited long-term clinical results have confirmed the safety and efficacy of this artificial disc.^{16,20} Most studies have been performed outside of North America, except for those associated with the ongoing United States Food and Drug Administration randomized prospective multicentre clinical trial of the CHARITÉ disc replacement versus lumbar fusion.²⁰ To our knowledge, ours is the first Canadian study that reports the results of lumbar TDA.

The purpose of our study is to prospectively report the clinical and radiographic results of the SB CHARITÉ III TDA performed at our Canadian centre with a minimum of 2 years of follow-up.

METHODS

Between April 2001 and November 2006, we prospectively followed 64 patients after they received a CHARITÉ artificial TDA. The primary indication for surgery was discogenic low-back pain. Inclusion and exclusion criteria were similar to those reported by previous studies.^{20,25} Inclusion criteria were a single-level symptomatic degenerative disc disease (DDD) on magnetic resonance imaging (MRI) that was confirmed by provocative positive discography and associated with persistent symptoms of primarily back and, in some cases, leg pain without overt nerve root compression, an Oswestry Disability Index

(ODI) score of 30 or more, a visual analogue scale (VAS) score of 40 or more and failure to respond to a minimum of 6 months of nonoperative treatment. In addition, the patient had to be able to tolerate an anterior abdominal approach.

Exclusion criteria were more than 3 mm of spondylolysis or retrolisthesis, osteoporosis, spinal deformity greater than 11°, 2 or more levels of DDD, facet joint arthrosis, previous thoracic or lumbar spine fusion and the presence of current or previous lower lumbar fractures.

All patients included in this study were informed of the option of disc arthroplasty or fusion. The risk and benefits of each procedure were thoroughly discussed, and they consented to be followed prospectively for the purpose of this study. Our hospital ethics review board approved the study protocol.

Operative technique

The surgical approach used was the transperitoneal approach, performed by a vascular surgeon. The surgical technique has been well described previously.²⁶ Particular attention was given to preparation of the end plates. Care was taken to achieve flat end-plate surfaces extending to the posterior aspect of the vertebral body to maximize bone metal contact area. A curette or burr was used to remove any posterior ridge or osteophytes that might limit appropriate posterior posturing of the prosthesis end plate during final insertion.

The patients were mobilized as tolerated on the first postoperative day and instructed to avoid lumbar extension beyond neutral for a period of 6 weeks postoperatively.

Outcome assessment

Outcome evaluation took place preoperatively and postoperatively at 3, 6 and 12 months and once a year thereafter. Validated subjective outcome measures included the ODI, VAS and Short Form-36 (SF-36) health survey version 2. We calculated the VAS separately for leg and low-back pain, assessing both pain frequency and pain intensity. We determined the SF-36 mental component summary score (MCS) and physical component summary score (PCS).

Preoperatively, we obtained an MRI scan and a discogram of the symptomatic degenerate disc to identify surgical candidates. Pre- and postoperative radiological assessment included standing anterior–posterior (AP) and lateral lumbar radiographs in the neutral position, extension and flexion. We used lateral radiographs to assess angle of sagittal rotation (AOR), intervertebral translation of the cephalad on the caudal vertebra, anterior vertical motion (AVM) of the prosthesis and posterior vertical motion (PVM) of the prosthesis. The AOR measured total motion between flexion and extension radiographs by

determining the angle subtended between perpendiculars to the superior and inferior end plates of the replaced disc space. The AVM represented the difference between flexion and extension in the vertical distance between the anterior edge of the superior and inferior prosthetic end plates. We calculated the PVM in a similar manner using the posterior edges of the superior and inferior prosthetic end plates to determine the difference in distance between flexion and extension. We used the Cobb technique to determine the lumbar lordosis using the superior end plate of both L1 and S1. We used a T_2 midsagittal MRI scan to measure the preoperative anterior and posterior disc heights. We measured the postoperative anterior and posterior disc height using digitized radiographs of the lumbar spine in the neutral position.

We compared postoperative radiographs with preoperative radiographs to detect evidence of adjacent level degeneration above and below the TDA, based on disc space height loss, sclerosis and osteophyte formation.




Fig. 1. Measurements of subsidence and radiographic disc size.

Table 1. Demographic and clinical characteristics at baseline of patients who received a CHARITÉ III artificial total disc arthroplasty

Characteristic	Mean (SD) [range]*
Sex, no. (%)	
Male	25 (39)
Female	39 (61)
Age, yr	38.9 (7.7) [21–59]
Body mass index	25.7 (3.2) [19.2–32.3]
Duration of back pain, mo	69 (60.9) [9–324]
Length of stay in hospital, d	4.7 (1.5) [2.2–11.3]
Work status, no. (%)	
Full- or part-time	53 (82.8)
Compensation	11 (17.2)
Smoker, no. (%)	35 (54.7)
Symptomatic level, no. (%)	
L4–L5	5 (7.8)
L5–S1	59 (92.2)

SD = standard deviation.
*Unless otherwise indicated.

In addition to the above measurements, we noted the subsidence of the metal component of the artificial disc. Subsidence describes the depth of migration of the metal component of the artificial disc into the vertebra body, and this depth was measured as the distance between the posterior part of the metal component of the prosthetic disc that migrated into the vertebra body and the surface of the end plate of the vertebra (Fig. 1).

All measurements were performed using Centricity Picture Archiving & Communications System, version 1.0 (GE Medical Systems). The ratio of magnification was calculated by dividing the actual anterior–posterior artificial disc dimension with the radiographic dimension measured on a lateral radiograph. This magnification ratio was then used to correct the magnification error in all the other measurements. Two of us (M.K. and K.I.) obtained the measurements. Intra- and interobserver reliability was assessed using the intraclass correlation (ICC).

RESULTS

Patients

Sixty-four consecutive patients enrolled in this prospective study: 39 women and 25 men. The mean age at time of surgery was 39 (standard deviation [SD] 7.7) years. Seven (11%) patients were lost to follow-up, leaving 57 patients included in our analysis. The mean follow-up was 55 (range 24–84) months. The demographic and clinical characteristics of patients are provided in Table 1. Table 2 describes the number of patients reviewed at each follow-up.

The mean duration of symptoms was 69 (range 9–240) months. Preoperatively, 43 of 57 (75%) patients reported isolated low-back pain, whereas the remaining 14 (25%) reported a combination of back pain and leg pain. Fifty-five percent of patients smoked at the time of the procedure. Table 3 illustrates the difference in preoperative outcome measures between the smoking and nonsmoking cohorts. The mean time off of work was 9.9 (SD 23.3) months. Eleven percent claimed workers' compensation

Table 2. Number of patients reviewed at each follow-up

Follow-up	No. (%)
Preoperative	64 (100)
6 wk	64 (100)
3 mo	63 (98.43)
6 mo	62 (96.87)
1 yr	57 (89.06)
2 yr	60 (93.75)
3 yr	55 (85.93)
4 yr	42 (65.62)
5 yr	33 (51.56)
6 yr	18 (28.12)
7 yr	11 (17.18)

status. Seven patients had a previous discectomy, 6 of these were performed at L5–S1.

There were 53 TDAs carried out at the L5–S1 level and 4 at the L4–L5. The mean intraoperative blood loss was 220 (range 50–700) mL. The mean postoperative hospital stay was 4.7 (range 2–8) days.

Clinical outcome measures

All outcome measures, including the ODI, VAS, MCS and PCS, showed postoperative improvement at the 3-month follow-up ($p < 0.001$), which persisted during the remaining follow-up period (Figs. 2–5).

Complications

There were 3 (4.7%) early complications: 1 patient experienced a superficial abdominal hematoma, which was treated conservatively, 1 experienced a retroperitoneal hematoma that required exploration and evacuation and 1 of the 27 male patients presented with retrograde ejaculation.

Two patients required revision surgery, representing a late complication rate of 3.1%. The first was a young competitive field hockey player who underwent an L5–S1 disc replacement. This patient, against medical advice, returned to playing field hockey only a few weeks postoperatively and sustained an L5 inferior end plate fracture following a fall. Conservative treatment failed to control her symptoms,

Measure	Smoking status, mean (SD)*		p value
	Smoker	Nonsmoker	
Preoperative no.	35	29	
SF-36			
Physical component score	30.5 (12.6)	35.6 (16.5)	0.20
Mental component score	42.2 (14.3)	53.6 (22.1)	0.027
Oswestry Disability Index score	52.2 (15.8)	49.2 (16.5)	0.49
Visual analogue scale score			
Intensity of back pain	7.9 (1.9)	8.1 (2.0)	0.64
Frequency of back pain	8.8 (1.9)	9.4 (1.0)	0.14
Intensity of leg pain	6.7 (3.2)	5.3 (3.1)	0.10
Frequency of leg pain	7.2 (3.1)	5.5 (3.4)	0.06

SD = standard deviation; SF-36 = Short Form-36 health survey.
*Unless otherwise indicated.




Fig. 2. Oswestry Disability Index scores of patients who received a CHARITÉ III artificial total disc arthroplasty over 84 months of follow-up.




Fig. 3. Visual analogue scale (VAS) scores for back pain and intensity among patients who received a CHARITÉ III artificial total disc arthroplasty over 84 months of follow-up.




Fig. 4. Visual analogue scale (VAS) scores for leg pain and intensity among patients who received a CHARITÉ III artificial total disc arthroplasty over 84 months of follow-up.




Fig. 5. Short Form-36 (SF-36) scores of patients who received a CHARITÉ III artificial total disc arthroplasty over 84 months of follow-up.

and she required a posterolateral instrumented fusion leaving the TDA in situ (Fig. 6). The second patient who underwent an L4-L5 disc arthroplasty was among the very first patients selected for disc replacement. In retrospect, preoperative MRI scans demonstrated that this patient had some evidence of adjacent level degeneration at the L5-S1 level and despite a technically adequate procedure at the L4-L5 level continued to have low-back pain. A L5-S1 posterolateral instrumented fusion was performed by another surgeon and therefore it is not possible to comment on the severity of her symptoms before the fusion procedure or on the exact location of the painful segment (replaced L4-5 or L5-S1). In total, the reoperation rate was 4.7% (3 patients).

Radiographic outcome

Table 4 reports the results of the motion assessment performed at the last follow-up. There was a statistically significant increase ($p < 0.001$) in the postoperative compared with preoperative lumbar lordosis (Fig. 7). The mean anterior disc height and posterior disc height increased by 2.4 times the preoperative height (Table 5). Subsidence




Fig. 6. L4-L5 fusion following an interior end plate fracture resulting in acute subsidence of total disc arthroplasty and severe low-back pain.

was present in 44 of 53 (83%) patients at the L5-S1 level and was exclusively seen at the posterior part of the inferior end plate of L5. The mean subsidence was 1.7 (range 0-4.8) mm, measured at 3 months after surgery. Although there was a marginal progression of subsidence beyond this time point, it was not statistically significant. Subsidence had no effect on the range of motion at the replaced segment and did not correlate with clinical outcome. In patients who underwent an L4-5 TDA, subsidence occurred at both end plates: mean subsidence was 0.9 mm at the inferior end plate of L4 and 1.75 mm at the superior end plate of L5. Comparison between preoperative and postoperative radiographs demonstrated no evidence of adjacent level disease. Reliability of the measurements was very high for both the intra- and interclass correlation ($ICC > 0.8$).

DISCUSSION

The clinical results demonstrate a satisfactory outcome in patients who underwent a TDA in our unit. Our results are

Table 4. Results of the radiographic measurements of motion at the replaced segment

Motion	Mean (range)
Sagittal rotation, °	6.5 (0.5-22.4)
Intervertebral translation, mm	1.1 (0-2.4)
Anterior vertical motion, mm	1.5 (0-4.8)
Posterior vertical motion, mm	1.5 (0.2-5.7)




Fig. 7. Mean preoperative and postoperative lumbar lordosis.

Table 5. Mean pre- and postoperative disc height

Time	Mean, mm	
	Anterior disc height	Posterior disc height
Postoperative	17.5	8.4
Preoperative	7.3	3.5
Postoperative → preoperative	10.2	5.0

similar to those previously published.^{16–22} As this was a prospective cohort study, no control group was available for comparison, so the similarity of the results to other published cohorts is important to recognize. The results of this cohort fulfill previously described criteria defining success after TDA or fusion, which included improvement in ODI of more than 25% at 24 months postoperatively, no device failure, no major complications and no neurologic deterioration.²⁰ The clinical outcome measures of the current study demonstrate a significant improvement of the ODI of about 50% postoperatively and a similar degree of improvement in the VAS and SF-36. In our series, there were no cases of device failure, no neurologic deterioration and no major complications. This significant improvement in the clinical outcome measures may not just be related to replacement of the degenerative painful disc, but also to unloading of the facet joints following restoration of the disc height and improvement of the lordosis. This theory has been supported by a prospective morphological study of facet joint integrity following a CHARITÉ disc replacement concluding that decreases in subchondral bone density of the facet joint at the replaced segment suggest a reduction in the loading of the posterior column following a disc replacement.²⁷ Furthermore, improvement in the disc height resulted in an increase in the foramen height, which can result in the relief of preoperative leg symptoms produced by degenerative foraminal stenosis.

The complication rate associated with TDA has been reported to be in the range of 1%–40%.^{25,18,28,29} The results of the current study demonstrate a relatively low complication rate of 4.7% for early and 3.1% for late complications, resulting in a reoperation rate of 4.7%. In a retrospective study of the CHARITÉ TDA, after a minimum 10 years the overall reoperation rate was 10.4%.¹⁶ The authors' view was that both the design and instrumentation changes as well as specific training for TDA procedures resulted in much shorter learning curves and ultimately fewer complications related to technical aspects of the procedure, particularly for surgeons already familiar with the anterior approach to the low lumbar spine.¹⁶ One of the 2 revision surgeries in our cohort was performed for persistent low-back pain, likely derived from an adjacent level to the arthroplasty. This reinforces the critical importance of patient selection.

The range in reported complications may be explained by differing surgical approach and technique. Following an anterior approach, 1 patient in our cohort experienced retrograde ejaculation; this compares favourably to a 10-fold increase in the rate of retrograde ejaculation reported using a transperitoneal approach.³¹ High rates of heterotopic ossification and spontaneous fusion have now been attributed to the surgical technique used at the time of implantation and particularly to the repair of the anterior longitudinal ligament (ALL) following the disc implantation.³³ Rates of complete ossification and spontaneous fusion have

ranged between 2.8% and 60%.^{19,32} In our cohort of patients, the ALL was not repaired, and there were no observed cases of heterotopic ossification.

Motion was maintained at the replaced segment. The mean 6.5° of sagittal rotation during flexion and extension compares favourably to the sagittal rotation reported in the literature,³¹ where at a 5-year follow-up the mean angle of sagittal rotation was 6.0° for the CHARITÉ group of patients compared with 1.0° for the fusion group. An interesting finding in our study was the variation in rotation for our patient cohort, which ranged between 0.5° and 22.4°. We believe this can be explained by a lack of standardization when performing the flexion and extension radiographs. We frequently noticed a substantial inconsistency in rotation among follow-up periods for particular patients. In these patients, despite the fluctuations in sagittal rotation among follow-up periods, clinical outcome measures remained consistent, supporting the concept that the variation in range was related more to patient effort than symptoms.

The mean intervertebral translation in our study was 1.1 mm, the AVM 1.5 mm and the PVM 1.5 mm, which fall within the physiologic range of translation previously described.¹⁴ It remains unknown what motion is required at one spinal segment to protect the adjacent segments from degeneration. It is theorized that the restoration or preservation of motion at the replaced segment can only be beneficial for the adjacent segment, provided that this motion follows a pattern close to the physiologic one. No radiographic evidence of adjacent level disease was seen in our cohort of patients; however, we recognize that our follow-up period was relatively short for this to be fully assessed. Subsidence of the metal prosthesis on the end plate was quite common without influencing the clinical outcome and the motion across the replaced segment.

CONCLUSION

The 2- to 7-year follow-up of this Canadian cohort of patients demonstrated satisfactory clinical and radiographic results in a carefully selected patient population. The radiographic assessment confirmed preservation and maintenance of motion of the replaced disc with a follow-up averaging 55 months.

Competing interests: None declared.

Contributors: Drs. S. Bailey and Gurr designed the study. Drs. Katsimihas, Issa, Rosas-Arellano and Gurr acquired the data, which they analyzed with Drs. Fleming and C. Bailey. Drs. Katsimihas, Issa and C. Bailey wrote the article, which Drs. S. Bailey, Fleming, Rosas-Arellano and Gurr reviewed. All authors approved publication of the article.

References

1. Möller H, Hedlund R. Surgery versus conservative management in adult isthmic spondylolisthesis — a prospective randomized study: part 1. *Spine (Phila Pa 1976)* 2000;25:1711–5.

2. Fritzell P, Hagg O, Wessberg P, et al.; Swedish Lumbar Spine Study Group. 2001 Volvo Award Winner in Clinical Studies: lumbar fusion versus nonsurgical treatment for chronic low back pain: a multicenter randomized controlled trial. *Spine* 2001;26:2521-32.
3. Fairbank J, Frost H, Wilson-MacDonald J, et al. Randomised controlled trial to compare surgical stabilization of the lumbar spine with an intensive rehabilitation programme for patients with chronic low back pain: the MRC spine stabilization trial. *BMJ* 2005;330:1233-8.
4. Bjarke Christensen F, Stender Hansen E, Laursen M, et al. Long-term functional outcome of pedicle screw instrumentation as a support for posterolateral spinal fusion. *Spine (Phila Pa 1976)* 2002;27:1269-77.
5. Burkus JK, Transfeldt EE, Kitchel SH, et al. Clinical and radiographic outcomes of anterior lumbar interbody fusion using recombinant human bone morphogenetic protein-2. *Spine (Phila Pa 1976)* 2002;27:2396-408.
6. Burkus JK, Heim SE, Gornet MF, et al. Is INFUSE bone graft superior to autograft bone? An integrated analysis of clinical trials using the LT-CAGE lumbar tapered fusion device. *J Spinal Disord Tech* 2003;16:113-22.
7. Park P, Gartron HJ, Gala VC, et al. Adjacent segment disease after lumbar or lumbosacral fusion: review of the literature. *Spine (Phila Pa 1976)* 2004;29:1938-44.
8. Ghiselli G, Wang JC, Bhatia NN, et al. Adjacent segment degeneration in the lumbar spine. *J Bone Joint Surg Am* 2004;86-A:1497-503.
9. Gillet P. The fate of adjacent motion segments after lumbar fusion. *J Spinal Disord Tech* 2003;16:338-45.
10. Katz V, Schofferman J, Reynolds J. The sacroiliac joint: a potential cause of pain after lumbar fusion to the sacrum. *J Spinal Disord Tech* 2003;16:96-9.
11. Manchikanti L, Boswell MV. Sacroiliac joint pain after lumbar fusion to the sacrum. *J Spinal Disord Tech* 2005;18(Suppl):S135.
12. Sihvonen T, Herno A, Paljäärvi L, et al. Local denervation atrophy of paraspinal muscles in postoperative failed back syndrome. *Spine (Phila Pa 1976)* 1993;18:575-81.
13. Cunningham BW, Gordon JD, Dmitriev AE, et al. Biomechanical evaluation of the total disc replacement arthroplasty: an in vitro human cadaveric model. *Spine (Phila Pa 1976)* 2003;28:S110-S117.
14. White AA, Panjabi MM. *Clinical biomechanics of the spine*. Philadelphia (PA): Lippincott Williams & Wilkins; 1990.
15. McAfee PC, Cunningham B, Holsapple G, et al. A prospective, randomized, multicenter Food and Drug Administration investigational device exemption study of lumbar total disc replacement with the CHARITÉ artificial disc versus lumbar fusion: part II: evaluation of radiographic outcome and correlation of surgical technique with clinical outcomes. *Spine (Phila Pa 1976)* 2005;30:1576-83.
16. David T. Long term results of one-level lumbar arthroplasty: minimum 10-year follow-up of the CHARITÉ artificial disc in 106 patients. *Spine (Phila Pa 1976)* 2007;32:661-6.
17. David T. Lumbar disc prosthesis: five year follow-up study on 66 patients. *J Bone Joint Surg Br* 1999;81:252.
18. Griffith SL, Shelokov AP, Büttner-Janzen K, et al. A multicenter retrospective study of the clinical results of the LINK SB Charité intervertebral prosthesis. The initial European experience. *Spine (Phila Pa 1976)* 1994;19:1842-9.
19. Zeegers WS, Bohnen LM, Laaper M, et al. Artificial disc replacement with the modular type SB Charité III: 2 year results in 50 prospectively studied patients. *Eur Spine J* 1999;8:210-7.
20. Blumenthal S, McAfee PC, Guyer RD, et al. A prospective, randomized, multicentre Food, and Drug Administration investigational device exemption study of lumbar total disc replacement with the CHARITÉ artificial disc versus lumbar fusion: part I: evaluation of clinical outcomes. *Spine (Phila Pa 1976)* 2005;30:1565-75.
21. Guyer RD, McAfee PC, Hochshuler SH, et al. Prospective randomized study of the Charité artificial disc: data from two investigational centres. *Spine J* 2004;4:252S-9S.
22. McAfee PC, Fedder IL, Saiedly S, et al. SB Charité disc replacement report of 60 prospective randomised cases in a United States Centre. *J Spinal Disord Tech* 2003;16:424-33.
23. Lemaire JP, Carrier H, Sariali el-H, et al. Clinical and radiological outcomes with the Charité artificial disc: a 10-year follow-up. *J Spinal Disord Tech* 2005;18:353-9.
24. Link HD. History, design and biomechanics of the LINK SB Charité artificial disc. *Eur Spine J* 2002;11(Suppl2):S98-105.
25. Bertagnoli R, Kumar S. Indications for full prosthetic disc arthroplasty: a correlation of clinical outcome against a variety of indications. *Eur Spine J* 2002;11(Suppl2):S131-6.
26. Geisler FH. Surgical technique of lumbar artificial disc replacement with the CHARITÉ artificial disc. *Neurosurgery* 2005;56:46-57.
27. Trouillier H, Kern P, Refior HJ, et al. A prospective morphologic study of facet joint integrity following intervertebral disc replacement with the Charité artificial disc. *Eur Spine J* 2006;15:174-82.
28. Cinotti G, David T, Prostacchini F. Results of disc prosthesis after a minimum follow-up period of 2 years. *Spine (Phila Pa 1976)* 1996;21:995-1000.
29. Lemaire JP, Skalli W, Lavaste F, et al. Intervertebral disc prosthesis: results and prospects for the year 2000. *Clin Orthop Relat Res* 1997;337:64-76.
30. Siepe CJ, Mayer MH, Heinz-Leisenheimer M., et al. Total lumbar disc replacement: different results for different levels. *Spine (Phila Pa 1976)* 2007;32:782-90.
31. Sasso RC, Kenneth BJ, LeHuec JC. Retrograde ejaculation after anterior lumbar interbody fusion: transperitoneal versus retroperitoneal exposure. *Spine (Phila Pa 1976)* 2003;28:1023-6.
32. Putzier M, Funk JF, Schneider SV, et al. Charité total disc replacement—clinical and radiographic results after an average follow-up of 17 years. *Eur Spine J* 2006;15:183-95.
33. Miyamoto S, Kuratsu S, Yonenobu K, et al. Evaluation of cell proliferating potentials in ossification of the spinal ligaments by argyrophilic nucleolar organizer region (AgNOR) staining. In: Kurokawa T, editor. *Annual Report of the Investigation Committee for Ossification of the Spinal Ligaments under the auspices of Japanese Ministry of Health and Welfare*. Tokyo (Japan): Department of Orthopedics, University of Tokyo; 1992. p. 160-5.
34. Guyer RD, McAfee PC, Banco RJ, et al. Prospective, randomized, multicenter Food, and Drug Administration investigational device exemption study of lumbar total disc replacement with the CHARITÉ artificial disc versus lumbar fusion: five-year follow-up. *Spine J* 2009;9:374-86.