

## A REVIEW OF TRAUMA SYSTEMS USING THE CALGARY MODEL

John Barry Kortbeek, MD

Surgeons caring for severely injured patients have witnessed tremendous change over the past 2 decades with the rapid evolution of trauma systems. This paper describes the evolution of trauma systems in Canada, using the one in Calgary as a model. Canadian system guidelines were produced by the Trauma Association of Canada in 1993. Participation in Canadian accreditation is accelerating as increasingly more centres across the country undergo external review each year. Reporting of trauma outcomes, including standardized mortality and a variety of performance measures, is becoming the norm. Injury is being treated as a disease with comprehensive control strategies aimed at reducing death and disability rates through prevention, treatment and rehabilitation.

Les chirurgiens qui traitent des blessés graves ont été témoins d'un changement énorme au cours des deux dernières décennies avec l'évolution rapide des systèmes de traitement des traumatismes. Cet article décrit l'évolution de ces systèmes au Canada à l'aide du modèle de Calgary. Les lignes directrices du système canadien ont été produites par l'Association canadienne de traumatologie en 1993. Au Canada, la participation à l'agrément des centres de traumatologie prend de l'ampleur : de plus en plus de centres d'un bout à l'autre du pays se soumettent à un examen de l'extérieur chaque année. La production de rapports sur l'issue du traitement des traumatismes, faisant état notamment des taux normalisés de mortalité et autres mesures de rendement, devient la norme. Les lésions sont traitées comme une maladie et des stratégies intégrées de prise en charge visent à réduire les taux de mortalité et d'incapacité par la prévention, le traitement et la réadaptation

Although the concepts of trauma care originate in antiquity, organized comprehensive civilian trauma systems are a relatively new phenomenon. The Royal College of Physicians and Surgeons of Canada launched the Trauma Association of Canada (TAC) in 1983. The Society was founded to provide a multidisciplinary vehicle for the promotion and development of trauma systems across Canada. During the late 1980s, some regional trauma centres received recognition

and funding from provincial ministries of health, principally in Ontario. In 1993, the TAC released its guidelines,<sup>1</sup> which were based upon the American College of Surgeons' document<sup>2</sup> but reflected the realities of Canadian trauma care, including a propensity for blunt trauma and greater geographic catchment areas. The guidelines also provided an inclusive model with categories for tertiary, district and rural trauma centres, while maintaining a minimum standard of commitment felt to be essential for

providing quality trauma care. Budget cutbacks and restructuring, due to government deficits and debt, precipitated regionalization of trauma services across Canada during the early 1990s. The effects of these changes are ongoing; however, they certainly have focussed attention and in some cases accelerated the development of regional trauma systems across the country. A voluntary system of accreditation has been promoted by the TAC and has seen active participation by many tertiary trauma facilities.

*From the Department of Surgery, Division of General Surgery and Division of Critical Care, University of Calgary, Calgary, Alta.*

*Part of the symposium "What's New in Trauma Care?" presented at the 66th annual meeting of the Royal College of Physicians and Surgeons of Canada, Vancouver, BC, Sept. 26, 1997.*

*Accepted for publication July 9, 1999.*

**Correspondence to:** Dr. John B. Kortbeek, Director, Regional Trauma Services, Calgary Regional Health Authority, Foothills Medical Centre, 1403 – 29th St. NW, Calgary AB T2N 2T9; kortbeek@supernet.ab.ca

© 2000 Canadian Medical Association (abstract and text/résumé)

**RESPONSIBILITY OF A TRAUMA CENTRE**

The responsibility of a tertiary trauma centre is not only to provide complete, coordinated and efficient care but to enhance the entire trauma system through a variety of activities. These include outreach activities, education, and injury prevention and control, as well as development of the data system, including registry and injury surveillance. The centre must also establish cooperative relationships, good communication, transport and bypass agreements with primary and secondary hospitals in the region to complement the entire trauma system.<sup>3</sup> A simplified schematic of a regional trauma system is illustrated in Fig. 1.

**INDEXES FOR MAJOR TRAUMA**

Major trauma has often been defined as being present in patients with an Injury Severity Score (ISS) greater than 16 whose risk of death was more than 10%, as defined by the major trauma outcome study.<sup>4</sup> Injury severity scoring, however, is retrospective, and major trauma presents as part of a continuum from minor to lethal injury. Studies of inclusive systems suggest that of all injuries presenting to hospital, 5% to 10% require the resources of a tertiary trauma centre.<sup>5,6</sup> In addition, the spectrum of injury extends from severe life-threatening polytrauma to outpatient injuries.<sup>7</sup> This is best illustrated by the injury pyramid (Fig. 2). The trauma system

accreditation guidelines, developed by the TAC in 1993, attempt to address the need for an inclusive model of a trauma system.

Identifying patients who potentially have multiple injuries consistent with major trauma and who would benefit from trauma treatment in a tertiary level trauma facility, has been a goal of the prehospital system. A variety of triage tools have been proposed and studied, including the Trauma Score<sup>8</sup> and CRAM's scale.<sup>9</sup> The prehospital index (PHI),<sup>10</sup> in conjunction with the mechanism of injury (MOI), was assessed by a prospective study in the Calgary region over a 6-month period, analysing more than 3000 emergency medical service (EMS) injury transports.<sup>11</sup> The combined score had 89% specificity and 78% sensitivity for detecting major trauma (Table I). The majority of patients in whom high ISSs were missed were the elderly who had falls from standing heights. The study concluded that PHI/MOI triage was effective but emphasized that any triage system should be tailored to the individual requirements of the community in which it is to be used.

**TRANSPORTATION OF TRAUMA PATIENTS**

Any evaluation of a Canadian trauma system must also reflect the reality that half of the patients presenting to the tertiary centre are transported from rural and regional centres, often covering great distances over many hours.<sup>12</sup> The quality of air ambulance transfers was assessed by van Wijngaarden and colleagues,<sup>13</sup> who demonstrated inappropriate care in less than 6% of cases and a very small overtriage rate of 5%. The study concluded that, given the difficulty of triage, the air ambulance system was likely underutilized. Many factors determine time from injury to tertiary care including pre-primary hospital




FIG. 1. Schematic diagram of a simplified regional trauma system.

time, time in the originating hospital before contacting dispatch and, finally, organization of the air ambulance transport system. Appropriate selection of patients who are physiologically unstable for rapid evacuation and transport also remains controversial.<sup>14</sup> These are all items of ongoing interest. Many centres have developed trauma hotlines or rapid access lines. The Calgary region now has a single 1-800 number for critically ill and injured patients, providing immediate access by rural physicians to the receiving tertiary trauma or critical care team, as well as the emergency flight physicians and the dispatch centre.

### MEASURES FOR EVALUATING TRAUMA CENTRES

Do trauma centres make a difference? A heterogeneous population, multiple health professionals and caregivers, and a variety of health organizations all are factors that challenge our ability to scientifically validate the effectiveness of trauma systems. Despite these challenges, a large body of evidence now supports the effectiveness of regionalized trauma systems.<sup>15-20</sup> The development of trauma registries has allowed the science of injury to progress.<sup>21</sup> These comprehensive databases have evolved to allow prospective data entry on severely injured patients. They support performance improvement through the review of cases identified by standard audit filters and death reviews. Research on outcomes and effectiveness, injury surveillance and support for allocation of resources are all benefits of the tool.<sup>22</sup> Analysis of the Trauma Injury Severity Score (TRISS), based on a major trauma outcome study, has been one method of survival analysis.<sup>23</sup> Most organized centres today report TRISS outcomes and *z* scores, demonstrating a significantly improved outcome compared with major trauma outcome study (MTOS)

data. The *z* score shows whether there is a significant difference between the centre's overall mortality and survival statistics and MTOS data.<sup>24</sup> The *W* statistic indicates the actual increased or decreased number of survivors per 100 trauma admissions. The *M* statistic is used to demonstrate whether the populations match.

TRISS analysis is limited, however, because it is a rather crude measure of outcome, and many patients who present to trauma centres that have registries are already intubated and initial physiologic data required for TRISS is not available.<sup>25</sup> These patients often represent the more seriously injured ones with higher mortality. Other techniques used to examine the effectiveness of particular trauma systems include internal and external death and morbidity reviews that study preventable and potentially preventable deaths and complications. Finally, examination of specific injury cohorts has been an effective means of the examining outcome. Smith and colleagues<sup>26</sup> demonstrated a significant reduction in death and complication rates in patients treated at trauma centres, compared with nontrauma centres, based on discharge abstract data and patient management categories. Nakayama and colleagues<sup>27</sup> examined outcomes of childhood trauma in the Pennsylvania trauma system. On the basis of TRISS analysis, they reported higher pediatric death rates in rural trauma centres. Cales,<sup>28</sup> in a landmark article in 1984, demonstrated significant improve-

ments in potentially preventable morbidity and mortality after the introduction of organized trauma care in Orange County. In 1995, Stewart and colleagues<sup>29</sup> demonstrated a significant increase in survival of 6% after Ontario Ministry of Health designation. Finally, a variety of performance indicators measure the effectiveness of trauma systems (e.g., prehospital time, time to laparotomy). These are now appearing in the public domain, with some systems publishing their results on their Web sites (e.g., [www.trauma.org](http://www.trauma.org), a British trauma site with links to many trauma centres).

Canadian evaluation of trauma systems is ongoing with attention to defining essential elements. Two important publications have originated from changes in the Quebec system. Sampalis and colleagues<sup>30</sup> demonstrated the effectiveness of regional bypass with destination policies in urban trauma systems. They then presented the results of an ambitious project analysing the effects of regionalization and the development




FIG. 2. The injury pyramid.

Table I

Effectiveness of the Triage Tool Used to Treat Patients Admitted With Blunt Trauma\*

Triage tool	Sensitivity, %	PPV, %	Specificity, %	NPV, %	Accuracy, %
Prehospital index alone	41	40	98	98	97
Mechanism of injury alone	73	18	91	99	90
Combined score	78	17	89	99	89

PPV = positive predictive value, NPV = negative predictive value.

\*Reproduced with permission from Bond RJ, Kortbeek JB, Preshaw RM. Field trauma triage: combining mechanism of injury with the prehospital index for an improved trauma triage tool. *J Trauma* 1997;43(2):282-7.

of a trauma care network from 1992 to 1998.<sup>31</sup> The death rate in major trauma patients (ISS greater than 12) fell from 52% to 18%. The large study population also permitted assessment of trauma system components and their relative impact, including pre-hospital time, trauma centre designation and transfer versus direct transport. In 1996, a trauma program with a dedicated clinical trauma service was introduced in Vancouver.<sup>32</sup> Important reductions in delays to the operating room, length of stay in the trauma unit and TRISS mortality resulted.

## INJURY CONTROL

Injury control is a term increasingly used to reflect a comprehensive approach to the disease of injury.<sup>33</sup> It represents a systems approach, including injury prevention, education and treatment through to rehabilitation and reintegration. The cure for injury is prevention. Injury continues to be the leading cause of death in the first 4 decades of life, and challenges cancer and heart disease for overall potential years of life lost.<sup>34</sup> Per capita, fatalities due to motor vehicle collisions have been falling in Western countries since 1965 (Fig. 3). This presumably is a result of improved vehicle safety

design, traffic regulation and enhanced trauma systems, as well as general awareness and education.<sup>35</sup> Further reductions in deaths due to motor vehicle accidents are obtainable as evidenced by the Victoria, Australia, casualty crash initiative.<sup>36</sup> This effort achieved a mortality reduction of almost 50% between 1990 and 1992. Further reductions are possible but only through heightened public awareness and education, stricter enforcement and effective lobbying for changes in legislation. Health restructuring and cutbacks have focussed attention on injury control and have seen developments across the country.

Alberta has recently funded a provincial office of injury control. The objectives of the centre are to promote comprehensive solutions including enhanced injury surveillance, adoption of TAC guidelines, changes in public awareness, enforcement and legislation. Similar programs are evolving across North America.<sup>37-39</sup> In addition, 1997 saw the launch of the National Trauma Registry by the Canadian Institute for Health Information in conjunction with the TAC.<sup>40</sup> This important project will support national injury surveillance as well as population-based research on injury and trauma. In 1999, a pro-

posal to link TRISS results from Canadian trauma centres and develop a national benchmark, CTRISS, was presented at the TAC meeting in Montreal. Despite these improvements, research dollars are also necessary to drive evidenced-based improvements in the system. Injury research remains grossly underfunded relative to other health care problems, particularly in view of its significance as a public health problem.<sup>41</sup>

## SUMMARY

Trauma system development has made great strides. Trauma systems and injury control remain heterogeneous across the country. National guidelines and associations represent a powerful tool for committed individuals and organizations to ensure that quality is not compromised and that this important problem is addressed. This is particularly relevant during a time of rapid change and fiscal restraint during which health care traditions are continually challenged. The Ottawa-Carleton system, applying a model assessing cost per quality-of-life year gained, has confirmed the cost-effectiveness of trauma systems in Canada.<sup>42</sup> Trauma will continue to strike down, through loss of life and limb, many productive members of our society in the coming years. The value of investing time and energy in enhanced injury control is apparent.

## References

1. Trauma Association of Canada. *Trauma system accreditation guidelines*. The Association; 1993
2. American College of Surgeons, Committee on Trauma. *Resources for optimal care of the injured patient*. Chicago: The College; 1993.
3. West JG, Williams MJ, Trunkey DD, Wolfert CC Jr. Trauma systems: current status — future challenges. *JAMA* 1988;259(24):3597-600.




FIG. 3. Comparison of the number of fatalities per 100 000 motor vehicles in Canada (diamonds), the United States (squares), the United Kingdom (triangles) and Australia (crosses) for 1960, 1970 and 1992.

4. Champion HR, Copes WS, Sacco WJ, Lawnick MM, Keast SL, Bain LW Jr, et al. The major trauma outcome study: establishing rational norms for trauma care. *J Trauma* 1990;30(11):1356-65.
5. Moore EE, Mattox KL, Feliciano DV. Trauma. 2nd ed. Norwalk (Conn): Appleton & Lange; 1991.
6. Ivatury RR, Cayten CG. *Penetrating trauma*. Baltimore: Williams & Wilkins; 1996.
7. Report from the Calgary Regional Health Authority Regional Injury Program, April 1997.
8. Moreau M, Gainer PS, Champion H, Sacco WJ. Application of the trauma score in the prehospital setting. *Ann Emerg Med* 1985;14(11):1049-54.
9. Gormican SP. CRAMS Scale: field triage of trauma victims. *Ann Emerg Med* 1982;11(3):132-5.
10. Koehler JJ, Baer LJ, Malafa SA, Meindertsma MS, Navitskas NR, Huizenga JE. Prehospital Index: a scoring system for field triage of trauma victims. *Ann Emerg Med* 1986;15(2):178-82.
11. Bond RJ, Kortbeek JB, Preshaw RM. Field trauma triage: combining mechanism of injury with the prehospital index for an improved trauma triage tool. *J Trauma* 1997;43(2):283-7.
12. Goldstein L, Doig C, Gant P, Powell G, Nijssen-Jordan C, Johnston R, et al. Analysis of rural to tertiary care patient transportation systems: an analytical tool for systematic review of time variables with different modes of transportation [abstract]. *J Trauma* 1997;43(2):394.
13. van Wijngaarden M, Kortbeek JB, Lafrenière R, Cunningham R, Joughin E, Yim R. Air ambulance trauma transport: a quality review. *J Trauma* 1996;41(1):26-31.
14. Cunningham P, Rutledge R, Baker CC, Clancy TV. A comparison of the association of helicopter and ground ambulance transport with the outcome of injury in trauma patients transported from the scene. *J Trauma* 1997;43(6):940-6.
15. Mullins RJ, Veum-Stone J, Hedges JR, Zimmer-Gembeck MJ, Mann NC, Southard PA, et al. Influence of a statewide trauma system on location of hospitalization and outcome of injured patients. *J Trauma* 1996;40(4):536-46.
16. Bazzoli GJ, Madura KJ, Cooper GF, MacKenzie EJ, Maier RV. Progress in the development of trauma systems in the United States. *JAMA* 1995;273(5):395-401.
17. Cales RH, Trunkey DD. Preventable trauma death: a review of trauma care systems development. *JAMA* 1985;254(8):1059-63.
18. Esposito TJ, Sanddal ND, Hansen JD, Reynolds S. Analysis of preventable trauma deaths and inappropriate care in a rural state. *J Trauma* 1995;39(5):955-62.
19. Shackford SR, Hollingworth-Fridlund P, Cooper GF, Eastman AB. The effect of regionalization upon the quality of trauma care as assessed by a concurrent audit before and after institution of a trauma system: a preliminary report. *J Trauma* 1986;26(9):812-20.
20. West JG, Trunkey DD, Lim RC. Systems of trauma care: a study of two counties. *Arch Surg* 1979;114:455-60.
21. Vestrup JA. Update on trauma registries and trauma scoring. *Can J Surg* 1990;33(6):461-3.
22. Rutledge R. The goals, development and use of trauma registries and trauma data sources in decision making in injury. *Surg Clin North Am* 1995;5(2):305-26.
23. Boyd CR, Tolson MA, Copes WS. Evaluating trauma care: the TRISS method. *J Trauma* 1987;27(4):370-8.
24. Cottingham EM, Shuffleberger CM, Townsend R. The power of the Z statistic: implications for trauma research and quality assurance review. *J Trauma* 1989;29(11):1500-9.
25. Rutledge R, Osler T, Emery S, Kromhout-Schiro S. The end of the Injury Severity Score (ISS) and the Trauma and Injury Severity Score (TRISS): ICISS, an International Classification of Diseases, ninth revision-based prediction tool, outperforms both ISS and TRISS as predictors of trauma patient survival, hospital charges, and hospital length of stay. *J Trauma* 1998;44(1):41-9.
26. Smith JS, Martin LF, Young WW, Macioce DP. Do trauma centers improve outcome over non-trauma centers: the evaluation of regional trauma care using discharge abstract data and patient management categories. *J Trauma* 1990;30(12):1533-8.
27. Nakayama DK, Copes WS, Sacco W. Differences in trauma care among pediatric and nonpediatric trauma centers. *J Pediatr Surg* 1992;27(4):427-31.
28. Cales RH. Trauma mortality in Orange County: the effect of implementation of a regional trauma system. *Ann Emerg Med* 1984;13(1):15-24.
29. Stewart TC, Lane PL, Stefanits T. An evaluation of patient outcomes before and after trauma center designation using Trauma and Injury Severity Score analysis. *J Trauma* 1995;39(6):1036-40.
30. Sampalis JS, Denis R, Frechette R, Brown R, Fleischer D, Mulder D. Direct transport to tertiary trauma centers versus transfer from lower level facilities: impact on mortality among patients with major trauma. *J Trauma* 1997;43(2):288-96.
31. Sampalis JS, Denis R, Lavoie A, Frechette P, Boukas S, Nikalis A, et al. Trauma care regionalization: a process-outcome evaluation. *J Trauma* 1999;46(4):565-81.
32. Simons R, Eliopoulos V, Laflamme D, Brown DR. Impact on process of trauma care delivery 1 year after the introduction of a trauma program in a provincial trauma center. *J Trauma* 1999;46(5):811-6.
33. Avery JG. Accident prevention — injury control — injury prevention — or whatever? *Inj Prev* 1995;1(1):10-1.
34. Wilkins K. Causes of death: how the sexes differ. *Health Rep* 1995;7(2):33-43.
35. Powles JW, Gifford S. Health of nations: lessons from Victoria, Australia. *BMJ* 1993;306:125-7.
36. Vulcan P. Analysis of reductions in Victorian road casualties, 1989 to 1992. Presented at the Kananaskis Injury Prevention Conference; 1996.
37. Rivara FP, Grossman DC, Cummings P. Injury prevention. First of two parts. *N Engl J Med* 1997;337(8):543-8.

38. Rivara FP, Grossman DC, Cummings P. Injury prevention. Second of two parts. *N Engl J Med* 1997;337(9):613-8.
39. National Committee for Injury Prevention and Control. Injury prevention: meeting the challenge. *Am J Prev Med* 1989;5(3 Suppl):1-303.
40. Canadian Institute for Health Information & The Trauma Association of Canada, National Trauma Registry minimum data set. Presented to the Trauma Registry Committee at the annual meeting of the Royal College of Physicians and Surgeons of Canada, Vancouver, September 1997.
41. Sampalis J. Presentation of the TAC Research Committee to the annual business meeting of the Trauma Association of Canada, Vancouver, September 1997.
42. Seguin J, Garber BG, Coyle D, Hebert PC. An economic evaluation of trauma care in a Canadian level trauma hospital. *J Trauma* 1999;47(3 Suppl): S99-103.

## SESAP Critique / Critique SESAP

### CATEGORY 11, ITEMS 34 AND 35

The CT in the present case shows a single large apical lesion with an apparent thick-walled cavity. Although the lesion pictured might be malignant, the most likely diagnosis is chronic granulomatous infection. Video-assisted thoracoscopy (VATS) might provide a diagnosis, but the lesion is too large to remove by thoracoscopy. VATS would therefore need to be combined with needle aspiration or needle biopsy, and in that case offers little advantage over transthoracic needle aspiration. Thoracotomy and resection is more aggressive than would be warranted for an infectious disease, and would be dangerous if he has, for example, active tuberculosis. Transthoracic needle aspiration under x-ray control would be relatively simple and safe, and would be the least invasive procedure that is likely to produce a diagnosis. Thoracentesis could be done, but there is only a small amount of pleural fluid present, and it would be unlikely to be diagnostic. Repeat bronchoscopy is not indicated, because it has already failed to produce a diagnosis.

The differential diagnosis includes a number of possibilities. Wegener's granulomatosis typically produces multiple nodules, not large cavitory lesions. Histoplasmosis may present with a solitary mass, but would be much less likely to cavitate; further, histoplasmosis does not typically present with such a large lesion as seen here. The positive skin test is of little value in areas where histoplasmosis is endemic, such as northwest Missouri, where most of the population is histoplasmin positive. It does rule out anergy, so the negative tuberculin test is important. Tuberculosis, while always a possibility, seems less likely for three reasons; although it can produce apical cavitory lesions, the cavities tend to be more thin-walled; it does not typically grow into the chest wall, as this lesion appears to do; and the tuberculin skin test is reliable if one can rule out anergy. Aspergillosis is classically a secondary invader in a tuberculosis cavity. It is seen as a primary pathogen only in patients who are immunocompromised, which this patient was not. Actinomycosis, by contrast, can produce a large lesion, often attached to the chest wall, with small and irregular cavitory areas. There may be invasion of the chest wall, and even fistula formation to the skin of the chest. Like tuberculosis, it can attack otherwise healthy individuals. The characteristic pathologic finding of actinomycosis is the finding of yellow "sulfur granules" within the lesion.

Actinomycosis is caused by *Actinomyces israelii*, a bacterium normally found in the oral cavity, in the tonsillar crypts, and around the gums. It produces a filamentous growth pattern. This microaerophilic bacterium is notoriously difficult to isolate; anaerobic culture conditions are required. It is sensitive to many antibiotics, and penicillin remains the drug of choice. Other antibiotics such as chloramphenicol, clindamycin, tetracycline, and erythromycin can be used if penicillin fails. Treatment is begun with intravenous penicillin for several weeks, or until clinical response is noted, and then continued with oral penicillin for two to three months. Antifungal drugs such as amphotericin B and fluconazole are not indicated. Despite the somewhat fungal-sounding name, actinomycosis is a bacterial infection. Surgical therapy is usually not necessary, but resection is sometimes done for a destroyed lobe, chronic abscess, or bronchiectasis. It should be delayed until adequate antibiotic treatment has been well begun. Empyema is sometimes seen, but can usually be adequately treated with tube thoracostomy.

34  C 35  E

#### References

1. Hsieh MJ, Liu HP, Chang JP, Chang CH: Thoracic actinomycosis. *Chest* 104:366-370, 1993
2. Robinson LA, Reed EC, Galbraith TA, et al: Pulmonary resection for invasive *Aspergillus* infections in immunocompromised patients. *J Thorac Cardiovasc Surg* 109:1182-1197, 1995