

Soft-tissue images. Portal vein thrombosis and gas formation: unusual presentation of colon cancer

A 61-year-old woman presented to the Emergency Department with severe right lower quadrant pain and sepsis. Investigations revealed microcytic anemia and a significantly elevated leukocyte count. Emergent computed tomography (Figs. 1 to 3) demonstrated a diffusely thickened ascending colon with nodular mucosa and portal vein thrombus extending into the superior mesenteric vein with gas formation, indicating perforation into the portal venous system. Blood cultures were positive for *Clostridium* spp.

At laparotomy, omental, peritoneal and liver metastases were found and there was gross perforation of the tumour into the mesentery, indicating the source of her portal sepsis. There was no evidence of intestinal ischemia. A hemicolectomy was done. Pathological examination of the resected specimen revealed a moderately differentiated adenocarci-




FIG. 1.



FIG. 2.

Section editors: David P. Girvan, MD,
and Nis Schmidt, MD

Submitted by Dennis Hong, MD, and
Peter Lovrics, MD, Department of Surgery,
McMaster University, Hamilton, Ont.

Submissions to *Surgical Images*, soft-
tissue section, should be sent to Dr.
David P. Girvan, Victoria Hospital
Corporation, PO Box 5375, Station B,
London ON N6A 5A5 or to Dr. Nis
Schmidt, Department of Surgery, St.
Paul's Hospital, 1081 Burrard St.,
Vancouver BC V6Z 1Y6

Correspondence to: Dr. Peter Lovrics,
Department of Surgery, St. Joseph's
Hospital, 50 Charlton Ave. E, Hamilton
ON L8N 4A6; lovricsp@mcmaster.ca

© 2001 Canadian Medical Association

noma of the colon. Postoperative management included anticoagulation and broad-spectrum antibiotics, including penicillin. She was discharged 2 weeks postoperatively and was taking coumadin. She died 3 months later of her disease.

Portal vein thrombosis is a rare occurrence with numerous causes, the

most common of which are cirrhosis and neoplastic disease in adults.¹ Anticoagulation is the treatment of choice for this condition when it is not associated with intestinal ischemia. Gas in the portal venous system is indicative of perforation of visceral origin and usually mandates immediate laparotomy. However, the pres-

ence of portal venous gas has been reported in various clinical settings, in many cases not requiring urgent laparotomy. An algorithm for the management of portal venous gas has been outlined by Hong and associates.² Similarly, clostridial sepsis is rare and is associated with a very high death rate despite advances in antibiotic therapy. It most often has a gastrointestinal source, either directly from gastrointestinal neoplasia or indirectly from mucosal translocation in a setting of weakened host resistance.³ Clostridial bacteremia is almost always associated with findings of sepsis and merits prompt investigation and treatment.



FIG. 3.

References

1. Cohen J, Edelman RR, Chopra S. Portal vein thrombosis: a review. *Am J Med* 1992;92:173-82.
2. Hong JJ, Gadaleta D, Rossi P, Esquivel J, Davis JM. Portal vein gas, a changing clinical entity. *Arch Surg* 1997;132:1071-5.
3. Myers G, Ngoi SS, Cennerazzo W, Harris L, DeCosse JJ. Clostridial septicemia in an urban hospital. *Surg Gynecol Obstet* 1992;174:291-6.



ASSOCIATION
MÉDICALE
CANADIENNE



CANADIAN
MEDICAL
ASSOCIATION

CANADIAN JOURNAL OF SURGERY

The *Canadian Journal of Surgery* is a peer-reviewed journal specific to the continuing medical education needs of Canada's surgical specialists. It is published bimonthly by the Canadian Medical Association and is available online at: www.cma.ca/cjs

CMA Member Service Centre

tel 888 855-2555 or 613 731-8610 x2307

fax 613 236-8864

cmamsc@cma.ca