

# Total hip arthroplasty in a patient with neurofibromatosis type I and recurrent spontaneous hip dislocation

Thierry Odent, MD;\* Pierre Ranger, MD;† Mehdi Aarabi, MD;‡ Reggie C. Hamdy, MD;‡ François Fassier, MD‡

A 26-year-old woman came to the hospital with her third spontaneous posterolateral dislocation of the left hip, which had occurred as this young woman was getting out of a boat (Fig. 1). Her history included a diagnosis at 9 months of age of neurofibromatosis (NF) type I.

Reduction of the hip dislocation with traction was easy under general anesthesia, with no resultant complications, as confirmed by standard radiographs, arthrogram and arthroskan. During the following 2 years she had 3 other episodes of hip dislocation from trivial trauma. A shelf operation and fascia lata tenorrhaphy led to a symptom-free period of 8 years until a dislocation recurred, with subsequent development of progressive hip arthritis.

She underwent total hip arthroplasty. During that procedure, we noted that her hip muscles were atrophied and her capsulo-ligamentous structures loose, but found was no evidence of neurofibroma in or around the hip joint. After the hip replacement, an intraoperative examination demonstrated poor stability. To tighten the gluteal muscles, we performed a trochanteric osteotomy, fixed with a Downmills hook.

Afterward the patient had another episode of dislocation and developed recurrent instability of her artificial joint. This necessitated the replacement of the acetabular component by a constrained liner (Fig. 2).

At follow-up 18 months later, she has had no further dislocations.

## Discussion

To the best of our knowledge, this is the first reported case of recurrent hip dislocation in a patient with NF without any evidence of intra-articular neurofibromas or bony anomalies of the hip joint. A review of the literature revealed only 7 hip dislocations that have been reported in

patients with NF I, including 2 gradual dislocations in 2 young children<sup>1</sup> and 5 acute dislocations after trivial trauma in older patients.<sup>2-5</sup>

The following triad was described in acute dislocations by Guillemet and colleagues<sup>2</sup>: skin lesions discovered at birth, hypertrophy of the lower limb in childhood, and spontaneous dislocation of the hip in adult life. All the cases of dislocations reported were associated with intra-articular neurofibromas and bony chan-



Fig. 1. Posterolateral hip dislocation (third episode).

The \*Service d'orthopédie et traumatologie infantile, Hôpital des enfants malades, Paris, France, the †Hôpital du Sacre-Coeur de Montréal and the ‡Shriners Hospital for Children, McGill University, Montréal, Que.

Accepted for publication Feb. 6, 2004

**Correspondence to:** Dr. François Fassier, Shriners Hospital for Children, 1529 Cedar Ave., Montréal QC H3G 1A6; fax 514 842-8664; ffassier@shrinenet.org

ges, including acetabular dysplasia, erosion and growth disturbance with resultant coxa valga caused by neurofibroma invasion. Furthermore, mechanical problems such as weakness of the abductor

muscles caused by spinal-cord tumours were also reported as possible predisposing factors.<sup>1</sup>

In our case, we found no anatomical or mechanical abnormalities such as neu-

rofibromas to explain the problem. The evolution toward degenerative changes, instability and dislocation could suggest a deficiency of the normal sensory protection of the hip. This hypothesis was emphasized by Phillips and McMaster<sup>5</sup> because of the absence of pain during progressive dislocation in neuropathic arthropathy.

For our patient, because her soft tissue was not strong enough to restrain the hip joint, further joint dislocation could be prevented only by mechanical stabilization, including a shelf operation and insertion of a constrained-liner acetabular component. Despite the short-term good result, long-term prognosis is of concern because of her high risk of loosening of the acetabular component and early wearing in the absence of normal sensation.

**Competing interests:** None declared.

#### References

1. Haga N, Nakamura S, Taniguchi K, Iwaya T. Pathologic dislocation of the hip in von Recklinghausen's disease: a report of two cases. *J Pediatr Orthop* 1994;14:674-6.
2. Guillemet M, Creyssel J, de Mourges G, Fischer L. Neurofibromatose de von Recklinghausen / Hypertrophie congénitale d'un membre inférieur dans l'enfance / Luxation spontanée de la hanche homolatérale. *Presse Med* 1970;78:1269-71.
3. Lachiewicz PF, Salvati EA, Hely D, Ghelman B. Pathological dislocation of the hip in neurofibromatosis. *J Bone Joint Surg Am* 1983;65:414-5.
4. Nakasone S, Norimatsu H, Hamasaki N, Kinjo Y, Ibaraki K, Yabiku. A case report of recurrent dislocation of the hip joint with neurofibromatosis. *Seikeigeka to Saigaijeka* 1989;38:511-4.
5. Phillips JE, McMaster MJ. Pathological dislocation of the hip in neurofibromatosis. *J R Coll Surg Edinb* 1987;32:180-2.



**Fig. 2.** Total hip arthroplasty with a constrained liner.