

Residents' quality of life during an orthopedic trauma rotation: a multicentre prospective observational study

Ali Zahrai, MD;* Mohit Bhandari, MD, MSc;† Amit Varma, MD;‡ William R. Rennie, MD, DABS;‡ Hans Kreder, MD, MPH;*§ David Stephen, MD;*§ Michael D. McKee, MD;*¶ James P. Waddell, MD;*¶ Emil H. Schemitsch, MD*¶

Objective: Fewer orthopedic surgery trainees are choosing careers in trauma. The impact of stressors on health-related quality of life has not been evaluated among orthopedic surgical residents. The current study aimed to determine predictors of unsatisfactory resident quality of life during trauma rotations. **Method:** This was a prospective observational study in which we chose orthopedic trauma rotations within 2 training programs at 3 level 1 trauma centres. A sample of 15 orthopedic surgery residents participated in the study. The main outcome measure was the 36-Item Short-Form Health Survey taken at baseline, 2 and 6 months into the rotation. Residents ranked level of stress for 15 work-related and 13 individual problems on 5-point Likert scales. **Results:** The residents ranged from training years 1 through 5, and their mean age was 31 (standard deviation [SD] 2.8) years. Fear of error decreased at 6 months, compared with baseline ($p = 0.007$). Lack of time for family decreased as a problem at 2 and 6 months, compared with baseline ($p = 0.025$). Baseline Physical Component Summary scores were above population norms ($p = 0.007$). At 2 months, scores for the concept role limitations (physical) decreased significantly ($p = 0.019$), compared with the baseline. Mean Physical Component Summary scores declined from 57.59 (SD 3.51) at baseline to 53.68 (SD 6.50) at 2 months ($p = 0.015$) and 53.94 (SD 3.52) at 6 months ($p = 0.008$). At 2 months, mean general health scores significantly decreased, compared with baseline ($p = 0.022$). Social functioning scores at 6 months were decreased from population norms ($p = 0.02$). More resident social time was spent with medical people exclusively at 6 months, compared with baseline ($p < 0.02$). **Conclusion:** Trauma rotations affect residents' emotional well-being. Training programs should equip residents with coping strategies to improve quality of life during trauma rotations.

Objectif : Moins d'apprenants en chirurgie orthopédique choisissent de faire carrière en traumatologie. On n'a pas évalué l'effet des facteurs de stress sur la qualité de vie reliée à la santé chez les résidents en chirurgie orthopédique. L'étude en cours visait à déterminer les prédicteurs de qualité de vie insatisfaisante chez les résidents au cours des stages en traumatologie. **Méthodes :** Au cours de cette étude prospective par observation, nous avons choisi des stages en traumatologie orthopédique organisés dans le cadre de 2 programmes de formation donnés à 3 centres de traumatologie de niveau 1. Un échantillon de 15 résidents en chirurgie orthopédique a participé à l'étude. La version abrégée d'un questionnaire sur la santé (le 36-Item Short-Form Health Survey) comportant 36 questions auxquelles les participants ont répondu au départ, et 2 et 6 mois après le début du stage, a constitué la principale mesure de résultats. Les résidents ont classé le niveau de stress relié à 15 problèmes professionnels et 13 problèmes individuels sur des

From the Divisions of Orthopedic Surgery, *Department of Surgery, University of Toronto, Toronto, and †Hamilton Health Sciences—General Hospital, Hamilton, Ont., the ‡Section of Orthopedic Surgery, Department of Surgery, Health Sciences Centre and University of Manitoba, Winnipeg, Man., §Sunnybrook and Women's College Health Sciences Centre and ¶St. Michael's Hospital, Toronto, Ont.

This paper was presented at the 11th Meeting of the Combined Orthopaedic Associations in Sydney, Australia, October 24–9, 2004, and at the Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS) in Chicago, Ill., March 22–6, 2006.

Accepted for publication June 16, 2007

Correspondence to: Dr. A. Zahrai, 100 College St., Rm. 302, Toronto ON M5G 1L5; ali.zahrai@utoronto.ca

échelles de Likert à 5 points. **Résultats :** Les résidents représentaient les années 1 à 5 de la résidence et avaient en moyenne 31 (écart-type [ET] 2,8) ans. La peur de l'erreur diminuait à 6 mois comparativement au début ($p = 0,007$). Le manque de temps à consacrer à la famille diminuait comme problème à 2 et 6 mois, comparativement au début ($p = 0,025$). Les résultats du sommaire des composantes physiques au début dépassaient les normes de la population ($p = 0,007$). À 2 mois, les résultats relatifs aux limitations du rôle (physique) diminuaient considérablement ($p = 0,019$), comparativement au niveau de référence. Les résultats moyens du sommaire des composantes physiques ont diminué pour passer de 57,59 (ET 3,51) au début à 53,68 (ET 6,50) à 2 mois ($p = 0,015$) et à 53,94 (ET 3,52) à 6 mois ($p = 0,008$). À 2 mois, les résultats moyens relatifs à l'état de santé général diminuaient considérablement par rapport au niveau de référence ($p = 0,022$). Les résultats liés au fonctionnement social à 6 mois n'atteignaient pas les normes de la population ($p = 0,02$). À 6 mois, les résidents passaient plus de temps en société avec des gens du milieu de la médecine exclusivement qu'au début ($p < 0,02$). **Conclusion :** Les stages en traumatologie ont une incidence sur le mieux-être affectif des résidents. Les programmes de formation devraient apprendre aux résidents des stratégies d'adaptation à suivre pour améliorer leur qualité de vie au cours des stages en traumatologie.

Quality of life and general well-being during orthopedic surgical residency is a topic of current interest, given evolving guidelines for maximum allowable work hours weekly.¹⁻⁴ The Accreditation Council for Graduate Medical Education (ACGME) requires all programs to limit resident work hours to 80 hours weekly, with some programs allotted an extra 10% for specific educational purposes.² Although physician fatigue has taken centre stage as the primary motivation behind this movement, the effect of these rules on the continuity of care for hospitalized patients needs to be critically analyzed from the perspectives of patients, physicians and the health care system.⁵

Damestoy and colleagues⁶ identified 3 main types of stressors in resident quality of life studies: work-related stressors such as long working hours, heavy workloads, time pressure and low self-confidence; personal stressors such as a lack of time for oneself and one's family; and economic stressors due to the financial burden of training. In a study by Lewittes and colleagues,⁷ 27% of nearly 2000 residents surveyed felt that they were "often" or "nearly always" bothered by the feeling that their workload was too heavy and could not be finished in a usual working day. Other studies have pointed out the link between excessive work demands, fatigue and quality of care.⁸⁻¹⁰ Residents' obligations to provide care while suffering from fa-

tigue may lead them to tolerate lower standards of patient care than they would if they were well rested.⁷ As well, fears of inadequate performance as a physician during residency may cause residents to ignore their personal well-being as they focus on their professional development.¹¹ Time pressures and sleep deprivation constitute the major stressors adversely affecting the ability of residents to learn and, at times, the quality of care they provide.¹¹⁻¹⁵

It is our belief that orthopedic surgery residents' quality of life is subject to adverse changes during a busy trauma rotation. Using this as our hypothesis, we chose the trauma rotation for the setting of our study. Trauma rotations of 3 or 6 months' duration are mandatory requirements of all orthopedic training programs in North America. The impact of stressors on health-related quality of life and well-being has not been thoroughly evaluated among orthopedic surgical residents on busy trauma rotations. The current study aimed to determine those modifiable factors and stressors that predict residents' quality of life. Further, information from this study might encourage trauma curriculum reform to renew enthusiasm for careers in trauma among orthopedic surgical trainees.

Methods

We conducted a prospective, multi-centre, observational study of quality

of life during standardized orthopedic trauma rotations across 3 level 1 trauma centres.

Eligibility criteria

Eligible trainees included orthopedic surgery residents rotating on an orthopedic trauma service during our study period. A total of 15 orthopedic surgery residents at 3 level 1 trauma hospitals (part of 2 orthopedic training programs) were enrolled in this study. Ten enrollees were senior residents (fourth and fifth year), and 5 enrollees were junior residents (first to third year).

Trauma rotations

Trauma rotations were at 3 level 1 trauma centres. The residents were on call an average of 1 in 4 days. The typical workday when not on call was from 0600 to 1800. The typical work week was 80 hours on average. Patient load varied depending on the service.

Baseline evaluation

Residents were approached before the start of their trauma rotation and asked to provide baseline information. This information included demographic characteristics of age, sex, year of residency, marital status and number of children. We measured overall stress, stress of being a resident and stress of combining a

personal and professional life (from Rudner and colleagues¹¹) before and during the rotation, using a 5-point Likert scale: 0 = little or no stress, 1 = minimal stress, 2 = moderate stress, 3 = high stress, 4 = excessive stress.

Assessment of quality of life

We assessed quality of life and general well-being with the 36-Item Short-Form Health Survey (SF-36),¹⁶ a stress questionnaire¹¹ and a list of individual and work-related problems experienced by residents.^{17,18}

On approval by the institutions' ethics review boards, survey questionnaires were given at baseline evaluation and at months 2 and 6 of the trauma rotation. The first survey served as a baseline, which we compared with subsequent surveys. Surveys required 20 minutes to complete. One research assistant analyzed all the surveys. Each resident was given a numeric code, and all identifiers were removed after analysis. Faculty and staff had no access to the surveys, and all results were kept anonymous and confidential.

The SF-36 includes 1 multi-item scale measuring each of the following 8 health concepts:

1. Physical functioning assesses limitations of normal physical activities and is designed to estimate the severity of limitation (10 questions).
2. Role limitations (physical) assesses functional limitations at work that are caused by physical health problems (4 questions).
3. Bodily pain assesses both the severity of pain and the extent to which it interferes with normal activities (2 questions).
4. Social functioning assesses the quantity and quality of interactions with others, extending measurement beyond the exclusively physical and mental health concepts (2 questions).
5. Mental health/emotional well-being assesses the 4 major mental health dimensions of anxiety, de-

pression, loss of behavioural or emotional control and psychological well-being (5 questions).

6. Role limitations (emotional) assesses functional limitations at work that are caused by emotional problems, but this concept excludes limitations caused by physical problems (3 questions).
7. Vitality/energy assesses a subjective feeling of well-being, including energy and fatigue (4 questions).
8. General health assesses physical health status and has been documented to be a good predictor of health care expenditures (5 questions).

We also derived Physical Component (physical functioning, role physical, bodily pain and general health perceptions) and Mental Component (vitality, social functioning, mental health and role emotional) Summary scores.¹⁹

In addition, we asked residents to rank 15 work-related stressors¹¹ on a 5-point Likert scale ranging from 0 (not a problem) to 4 (extreme problem). The 15 work-related stressors were as follows: learning everything, feeling rushed, being on call, self-doubt, fear of error, no time for needy patients, frustration, death of patient, conflict between patients, program problems, status as a resident, problems with other specialties, sleep problems, faculty problems and resident problems.

An inventory developed by Nelson and Harvey¹⁸ at the University of Minnesota, and a modified version developed by Smith, Andrasik and Quinn (revised scale),¹⁷ both having over 100 questions, yielded the top 10 individual problems identified by residents. Our surveys included these 10 problems as well as an additional 3 that we believed to be highly applicable to our study. Again, residents ranked the 13 individual problems on a 5-point Likert scale ranging from 0 (not a problem) to 4 (extreme problem). The 13 individual problems were as follows: not enough time for self, not enough

leisure time, not enough time for physical activities (sports, exercise), not enough time to spend with spouse/friends/relatives, problems created by inflexible work hours, not enough time for study, not confident enough about self in medicine, reservations about being a doctor, not enough money, sexual drive decreased since beginning residency, not sure about choice of residency, feeling unfairly treated by program staff, and residents or medical people the only people seen socially.

Statistical analysis

For each time period surveyed, the number of residents responding varied slightly. As a result, we used paired *t* tests to detect changes in the same group of residents throughout the rotations. Junior residents had shorter rotations and thus were not included in the 6-month follow-up. We compared each of the SF-36 domains with the mean normative SF-36 Canadian population domain scores²⁰ for people aged 25–34 years. Statistical significance was established with Student's *t* test to compare the means. We corrected our *p* values for multiple comparisons and conducted such tests to explore the specific points at which differences might have been important (or statistically significant). Because we carried out numerous *t* tests, we set statistical significance at a *p* value of less than 0.03 to avoid significance due to chance alone.

Results

In total, 15 residents (13 men and 2 women) from training years 1 through 5 participated in the study. Their mean age was 31 (standard deviation [SD] 2.8) years. Of the residents, 53.3% were married or in a common-law relationship, 46.7% were single, and 33.3% had children.

The overall stress level was ranked by residents at baseline and at 2 and 6 months. According to paired

t tests, there were no statistical differences in stress levels at baseline, 2 and 6 months. At 6 months, fear of error had decreased significantly from baseline ($p = 0.007$) (Table 1).

Throughout the rotation, there were no statistically significant changes in residents' perception of time for themselves. Lack of time for spouse, family and friends decreased as a problem among residents at 2 and 6 months, compared with baseline ($p = 0.025$) (Table 2). In other words, residents reported having more time to spend with their spouse, family and friends. At 6 months, more resident social time was spent with medical people exclusively, compared with baseline ($p = 0.018$).

For the SF-36, mean baseline role limitation (physical) scores of residents were 97.32 (SD 5.32). At 2 months, these scores significantly decreased to 83.04 (SD 24.2) ($p = 0.019$) when compared with the baseline score (Table 3, Fig. 1). There were no statistically significant differences among residents' scores at baseline, 2 and 6 months when

compared with the mean Canadian population norms for role limitation.

Residents' mean scores at baseline for bodily pain, vitality, mental health, physical and social functioning and role limitations (emotional) did not change significantly over the rotation. At 6 months, however, the residents' mean social functioning scores had decreased significantly from the Canadian norms (mean 86.3, SD 20.3; $p = 0.02$) (Fig. 2).

At baseline, mean scores for general health were 83.14 (SD 13.10). At 2 months, this score had decreased significantly to 76.64 (SD 14.07) ($p = 0.022$ v. baseline) (Table 3, Fig. 1). The mean 6-month general health score (75.78, SD 16.14) was also lower than baseline scores, but this was not statistically signi-

ficant ($p = 0.037$). No statistically significant differences were found for residents' mean scores at baseline, 2 and 6 months when compared with the mean Canadian population norms for general health.

Physical Component Summary (PCS) scores take into account scores for physical functioning, role (physical), bodily pain and general health. PCS scores declined from 58.13 (SD 2.93) at baseline to 53.68 (SD 6.50) at 2 months ($p = 0.015$), and 53.94 (SD 3.52) at 6 months ($p = 0.008$) (Table 3, Fig. 1). The baseline mean PCS score for the residents was significantly higher ($p = 0.007$) than the mean Canadian population PCS score (53.0, SD 7.2) (Fig. 2). No statistically significant differences were found among residents' mean scores at 2 and 6 months when compared with the mean Canadian population norms for the PCS.

The Mental Component Summary (MCS) score includes scores for vitality, social functioning, mental health and role (emotional). The mean MCS score for residents did not change over time, and there were no statistically significant differences between residents' mean scores at baseline, 2 and 6 months and the mean Canadian population norms for the MCS.

Discussion

There is no question that surgical residency programs impose an added stress on quality of life and well-being, mainly owing to the high volume of work-related experience that is required to train a successful surgeon. In various residency programs, including the surgical specialties, Perry and Osborne²¹ studied the following characteristics addressed before and after residency: eating habits, hours of sleep, recreational time, exercise and missed significant events. They found that residents perceived significant changes in many areas of health and wellness during residency training. In a survey

Table 1
Changes from baseline in residents ranking of work-related stressors

Stressor	Significant change	
	2 mo	6 mo
Learning everything	No	No
Feeling rushed	No	No
Being on call	No	No
Self-doubt	No	No
Fear of error	No	Yes*
No time for needy patient	No	No
Frustration	No	No
Death of patient	No	No
Conflict between patients	No	No
Program problems	No	No
Status as a resident	No	No
Problems with other specialties	No	No
Sleep problems	No	No
Faculty problems	No	No
Resident problems	No	No

*Decreased, $p = 0.007$.

Table 2
Changes from baseline in residents' ranking of individual problems

Problem	Significant change	
	2 mo	6 mo
Not enough time for self	No	No
Not enough leisure time	No	No
Not enough time for physical activities (exercise, sports)	No	No
Not enough time to spend with spouse, friends, relatives	Yes*	Yes*
Problems created by inflexible work hours	No	No
Not enough time for study	No	No
Not confident enough about self in medicine	No	No
Reservation about being a doctor	No	No
Not enough money	No	No
Sexual drive decreased since beginning residency	No	No
Not sure about choice of residency	No	No
Feeling treated unfairly by program staff	No	No
Residents or medical people only people seen socially	No	Yes†

*Decreased, $p = 0.025$
†Increased, $p = 0.018$

of residents throughout their first year, Bruce and colleagues²² found that interns experienced increased levels of stress and ill health in the middle of the year as compared with the start and end of the year.

In evaluating the trainees' quality of life during a busy trauma rotation, we used several modalities. The SF-36 general quality of life validated survey has not, to our knowledge, been used with surgical residents before this

study. The SF-36 has been validated across several disease states as well as "healthy norms." We simply applied this questionnaire to a population otherwise perceived as healthy and normal. This tool allowed us to compare the general health and emotional well-being of residents with age-matched population norms. Our findings suggest that differences do occur in this population and deserve future exploration in larger data sets.

The SF-36 showed that resident scores declined in 2 out of 8 health domains and in the PCS. General health scores declined significantly at 2 months, compared with baseline scores. PCS scores declined from mean 57.59 (SD 3.51) at baseline to mean 53.68 (SD 6.50) at 2 months ($p = 0.015$), and mean 53.94 (SD 3.52) at 6 months ($p = 0.008$). At 6 months, residents' social functioning scores had decreased significantly from the Canadian norms (mean 86.3, SD 20.3; $p = 0.02$). It has been shown that a 5-point difference between groups or a 5-point change over time is considered clinically relevant.²³ Given that the baseline physical function and PCS scores were above the Canadian population's mean, the decrease in the PCS score at 2 and 6 months is indicative of a decline in the residents' general health through the rotation.

Role limitations (physical) assess the individual's functional limitations at work that are caused by physical health problems, and at 2 months, resident scores in this domain decreased from baseline ($p = 0.019$). It is thus imperative for residents to be in good physical condition before starting their trauma rotation, as some decline is to be expected.

Fear of error decreased among residents at 6 months when compared with baseline ($p = 0.007$). It is evident from our study that, as the residents progressed in their rotation, their knowledge base and technical experience increased. Reduced fear of error was the most important stress-reducing factor modified during the rotation. Lack of time for spouse, family and friends decreased as a problem at 2 and 6 months when compared with baseline ($p = 0.025$). Therefore, issues that initially were significantly problematic were diminished or nonexistent as the rotation went on. This confirms that coping strategies, when placed early in the training program, can be beneficial and thus decrease the level of anxiety and stress among residents early in

Table 3

SF-36 scores at baseline compared with 2- and 6-month scores

SF-36 general health domains	Scoring period; mean score (and SD)		
	Baseline <i>n</i> = 14	2 months <i>n</i> = 14	6 months <i>n</i> = 9
Physical functioning	98.93 (2.89)	94.92 (8.38) $p = 0.048$	98.33 (2.50) $p = 0.04$
Role limitations (physical)	97.32 (5.32)	83.04 (24.2) $p = 0.019^*$	85.42 (21.65) $p = 0.066$
Bodily pain	85.64 (19.60)	76.21 (26.48) $p = 0.16$	74.44 (22.28) $p = 0.10$
Social functioning	83.04 (23.82)	84.82 (22.02) $p = 0.38$	72.22 (27.80) $p = 0.23$
Mental health	77.50 (14.51)	74.29 (16.39) $p = 0.19$	80.00 (16.77) $p = 0.084$
Role limitations (emotional)	85.71 (17.11)	86.31 (18.38) $p = 0.45$	82.41 (29.30) $p = 0.15$
Vitality	63.39 (16.60)	58.48 (17.78) $p = 0.22$	57.64 (18.16) $p = 0.053$
General health	83.14 (13.10)	76.64 (14.07) $p = 0.022^*$	75.78 (16.14) $p = 0.037$
PCS	58.13 (2.93)	53.68 (6.50) $p = 0.015^*$	53.94 (3.52) $p = 0.008^*$
MCS	47.89 (9.62)	48.27 (9.42) $p = 0.43$	47.10 (12.97) $p = 0.36$

MCS = mental component summary; PCS = physical component summary; SD = standard deviation; SF-36 = 36-Item Short-Form Health Survey.
*Paired *t* test statistically significant change at $p < 0.03$.




FIG. 1. SF-36 domains with significant changes. PCS = Physical Component Summary score; SF-36 = 36-Item Short-Form Health Survey.

their rotation. Rudner's recommendations,¹¹ based on survey results, include developing support groups, providing advisors and assigning a senior resident to each beginning first-year resident. In our training program, each staff surgeon mentors a single resident, and regular one-on-one meetings take place. We believe this is also highly valuable for career counselling and helps to support residents during difficult times.

A recent study by Sargent and colleagues²⁴ found considerable burnout in orthopedic surgery residents, with high emotional exhaustion and depersonalization. Protective factors included being a parent, having a physician father, spending time with spouse and discussing concerns with colleagues, friends and family. Schwartz and colleagues²⁵ found that "high-quality" teaching rounds and a night-float system were among the suggestions made by the residents to lessen stress. Further, to cope with stress, the residents reported that they talked to others, tried to see humour in the situation or slept. This was illustrated in our study, where more resident social time was spent exclusively with medical people at 6 months compared with baseline.

We realize that this study is limited by its small sample size. We also realize that this study is specific to orthopedic surgery residents, our target group. We acknowledge the limitations of this study specifically related to sample size. Whether our findings are real will only be revealed by evaluations conducted by other investigators in a similar subset of people. Ultimately, increasing the sample should improve the precision of our estimates and also provide improved generalizability. We do believe, however, that similar studies within surgical and nonsurgical subspecialties should take place in the near future. The importance of such studies for improving the surgical education and quality of life of residents cannot be underestimated.

Trauma rotations affect residents' emotional well-being. Training programs should equip residents with coping strategies to improve quality of life during trauma rotations. Mentorship programs wherein residents meet their mentors regularly have been implemented in some programs, including ours. Further, stress management training should be implemented and has been shown to be beneficial if frequent training is pro-

vided; otherwise, its benefits are not long-lasting.²⁶ Stress management programs may also be beneficial to residents before they start trauma rotations. Improving the quality of life for orthopedic residents in a trauma rotation should begin early in their training with support programs implemented by residency programs.

Competing interests: None declared.

Contributors: Drs. Zahrai, Bhandari, Rennie, Stephen and Schemitsch designed the study. Drs. Zahrai, Varma, Kreder and McKee acquired the data, which Drs. Zahrai, Waddell and Schemitsch analyzed. Drs. Zahrai and Bhandari wrote the article. All authors reviewed the article and provided final approval for publication.

References

- Gelfand DV, Podnos YD, Carmichael JC, et al. Effect of the 80-hour workweek on resident burnout. *Arch Surg* 2004;139:933-40.
- Cockerham WT, Cofer JB, Lewis PL, et al. Resident work hours: Can we meet the ACGME requirements? *Am Surg* 2004;70:687-90.
- Chandra RK. The resident 80-hour work week: How has it affected surgical specialties? *Laryngoscope* 2004;114:1394-8; discussion 1319.
- Mendoza KA, Mendoza B, Britt LD. A template for change and response to work hour restrictions. *Am J Surg* 2003;186:89-96.
- Fletcher KE, Saint S, Mangrulkar RS. Balancing continuity of care with residents' limited work hours: defining the implications. *Acad Med* 2005;80:39-43.
- Damestoy N, Brouillette L, De Courval LP. Stress and residents' lifestyle. *Can Fam Physician* 1993;39:1576-80.
- Lewittes LR, Marshall VW. Fatigue and concerns about quality of care among Ontario interns and residents. *Can Med Assoc J* 1989;140:21-4.
- Rosen IM, Bellini LM, Shea JA. Sleep behaviors and attitudes among internal medicine housestaff in a U.S. university-based residency program. *Acad Med* 2004;79:407-16.
- Papp KK, Stoller EP, Sage P, et al. The effects





FIG. 2. Significant change in SF-36 scores compared with population norms. PCS = Physical Component Summary; SF-36 = 36-Item Short-Form Health Survey.

- of sleep loss and fatigue on resident-physicians: a multi-institutional, mixed-method study. *Acad Med* 2004;79:394-406.
10. Parshuram CS, Dhanani S, Kirsh JA, et al. Fellowship training, workload, fatigue and physical stress: a prospective observational study. *CMAJ* 2004;170:965-70.
 11. Rudner HL. Stress in family practice residents. *Can Fam Physician* 1986;32:319-23.
 12. Gaba DM, Howard SK. Patient safety: fatigue among clinicians and the safety of patients. *N Engl J Med* 2002;347:1249-55.
 13. Leonard C, Fanning N, Attwood J, et al. The effect of fatigue, sleep deprivation and onerous working hours on the physical and mental wellbeing of pre-registration house officers. *Ir J Med Sci* 1998;167:22-5.
 14. Samkoff JS, Jacques CH. A review of studies concerning effects of sleep deprivation and fatigue on residents' performance. *Acad Med* 1991;66:687-93.
 15. Lewittes LR, Marshall VW. Fatigue and concerns about quality of care among Ontario interns and residents. *CMAJ* 1989;140:21-4.
 16. Ware JE, Sherbourne CD. The MOS 36-Item Short-Form Health Survey (SF-36). *Med Care* 1992;30:473-83.
 17. Smith MF, Andrasik F, Quinn SJ. Stressors and psychological symptoms of family practice residents and spouses. *J Med Educ* 1988;63:397-405.
 18. Nelson EG, Henry WF. Psychosocial factors seen as problems by family practice residents and their spouses. *J Fam Pract* 1978;6:581-9.
 19. Ware JE Jr, Kosinski M, Keller SD. *SF-36 physical and mental health summary scales: a user's manual*. Boston: The Health Institute, New England Medical Center; 1994.
 20. Hopman WM, Towheed T, Anastassiades T. Canadian normative data for the SF-36 health survey. *CMAJ* 2000;163:265-71.
 21. Perry MY, Osborne WE. Health and wellness in residents who matriculate into physician training programs. *Am J Obstet Gynecol* 2003;189:679-83.
 22. Bruce C, Thomas PS, Yates DH. Health and stress in Australian interns. *Intern Med J* 2003;33:392-5.
 23. Ware JE Jr. *SF-36 health survey manual and interpretation guide*. Boston: The Health Institute, New England Medical Center; 1993.
 24. Sargent CM, Sotile W, Sotile MO, et al. Stress and coping among orthopaedic surgery residents and faculty. *J Bone Joint Surg Am* 2004;86:1579-86.
 25. Schwartz AJ, Black ER, Goldstein MG, et al. Levels and causes of stress among residents. *J Med Educ* 1987;62:744-53.
 26. Rowe MM. Teaching health-care providers coping: results of a two-year study. *J Behav Med* 1999;22:511-27.



Coming Soon to the
**Canadian Journal
of Surgery**

Manuscript Central — an innovative, online submission and peer-review system for scholarly publishers

It automates manuscript submission, streamlines reviewing and revision and reduces time to decision!

Check in at <http://mc.manuscriptcentral.com/cjs> for a launch date in June