

# Breast cancer metastasizing to the tongue

Pratap Neelakantan, MD;\* Sean R. McLean, MD;\* Sinead Kenny, RCN;\* Myooran Nathan, MD;\* Lajja Panchal, MD;† Gurpreet Sandhu, MD;‡ Charles R. Coombes, MD;\* Carlo Palmieri, MD\*

Involvement of the oral cavity by metastatic disease is rare and can include either bone or soft tissue. In this note, we describe the case of a young woman who had a metastatic deposit on her tongue, secondary to primary breast cancer.

## Case report

A 24-year-old woman presented with a lump in the right breast. A right mastectomy and axillary clearance with immediate reconstruction was performed. The operative specimen revealed a 38-mm, grade 2, invasive ductal carcinoma that gave weakly positive results for estrogen receptors (ERs) but negative findings for progesterone receptors (PRs) and human epidermal growth factor receptor 2 (HER2). One of 14 axillary lymph nodes was found to contain cancer cells, and

there was extracapsular spread. She received 6 cycles of adjuvant 5-fluorouracil (600 mg/m<sup>2</sup>), epirubicin (60 mg/m<sup>2</sup>) and cyclophosphamide (600 mg/m<sup>2</sup>) followed by adjuvant radiotherapy and was subsequently started on tamoxifen. Ten months after she completed chemotherapy, she presented with shortness of breath on exertion and a painful ulcer on the lateral aspect of the tongue that had been present for the preceding 2 months. Examination revealed dullness and reduced breath sounds over the right lung base and an ulcerated area over the lateral aspect of the tongue, with extensive induration. Computed tomography revealed a right-sided pleural effusion and multiple parenchymal masses in the lung fields, consistent with metastatic disease. A biopsy of the tongue lesion revealed a high-grade invasive carcinoma that ap-

peared focally to be forming glands (Fig. 1, left); the tumour expressed cytokeratin 14 (Fig. 1B) and cytokeratin 7 but was negative for ER, PR and HER2. Comparison was made with the primary breast cancer, and the lesion was confirmed as a metastatic deposit from that tumour. Chemotherapy with docetaxel 100 mg/m<sup>2</sup> stabilized the tongue metastasis, but there was progression of the pulmonary metastasis. She received further chemotherapy but died from her disease a little over 2 years from the time of original presentation.

## Discussion

An oral metastatic deposit is the first manifestation of an occult primary malignant tumour in up to one-third of cases. A review of over 6000 autopsies revealed 12 cases of




FIG. 1. Tongue biopsy showing (A) high-grade anaplastic carcinoma (hematoxylin-eosin, original magnification  $\times 200$ ) and (B) immunostaining for cytokeratin 14 (original magnification  $\times 25$ ).

From the Departments of \*Medical Oncology, †Histopathology and ‡Maxillofacial Surgery, Charing Cross Hospital, London, UK.

Accepted for publication July 25, 2007

**Correspondence to:** Dr. C. Palmieri, Department of Medical Oncology, Charing Cross Hospital, Fulham Palace Rd., London W6 8RF, UK; fax 44 2088461433; c.palmieri@imperial.ac.uk

lingual metastases. The most common primary tumour was malignant melanoma (5 cases), followed by breast and lung cancer (2 cases each); other primary sites were the colon, pancreas and esophagus.<sup>1</sup> The base of the tongue was the most common anatomic location, the lesion being clinically apparent in 83% of cases, although the diagnosis was only made before death in one-third of cases, and in 1 case, the lingual metastasis was the only apparent site of metastatic disease.<sup>1</sup> Four cases have been reported in the English literature of metastatic deposits to the tongue secondary to breast cancer: 2 involved the mobile part,<sup>2,3</sup> 1 the tip<sup>4</sup> and 1 the base of the tongue.<sup>5</sup> In 1 case, the primary and metastatic lesions were diagnosed concurrently<sup>2</sup>; in another case, the patient had established metastatic

disease<sup>3</sup>; and in 2 cases, it was the first sign of recurrent disease.<sup>4,5</sup> Recurrences occurred between 14 and 29 months after the initial diagnosis.<sup>3-5</sup> There were other sites of metastatic involvement in all cases. Patients were treated with surgical excision,<sup>2,4</sup> external beam radiotherapy<sup>3-5</sup> and chemotherapy with thiotepa (1 case).<sup>2</sup> Two of the patients died within 4 months of presentation,<sup>2,5</sup> the outcome in 1 patient was not reported,<sup>4</sup> and the remaining patient was alive and disease free 6 months after the diagnosis.<sup>3</sup>

**Competing interests:** None declared.

### References

1. Zegarelli DJ, Tsukada Y, Pickren JW, et al. Metastatic tumour of the tongue. Report of twelve cases. *Oral Surg Oral Med Oral Pathol* 1973;35:202-11.
2. Meher-Homji DR, Gavadia ML, Dabhoiwala NF. Carcinoma of the breast metastatic to the anterior part of the tongue. *Indian J Cancer* 1967;4:340-2.
3. Perchick JS, Kim TH. Breast cancer metastatic to the tongue. *J Oral Maxillofac Surg* 1986;44:484-6.
4. Fink I, Garb J. Carcinoma of the tip of the tongue: a case of metastasis from a malignant tumour of the breast. *Am J Surg* 1943;63:138-41.
5. Kolson H. Adenocarcinoma of the breast metastatic to the base of the tongue. *Laryngoscope* 1966;76:531-6.