

Postoperative bedrest improves the alignment of thoracolumbar burst fractures treated with the AO spinal fixator

Yen Dang, MD*
David Yen, MD*
Wilma M. Hopman, MA[†]

From the Departments of *Surgery and
†Community Health and Epidemiology,
Queen's University, Kingston, Ont.

Presented in part at the annual meeting
of the Canadian Spine Society,
Mar. 22–24, 2001, Mont Tremblant, Que.

Accepted for publication
Feb. 2, 2008

Correspondence to:
Dr. D. Yen
Douglas 5, Kingston General Hospital
76 Stuart St.
Kingston ON K7L 2V7
fax 613 548-2518
yend@kgh.kari.net

Background: A loss of reduction due to inadequate support of the anterior column when using short-segment instrumentation to treat burst fracture and novel methods for support of the anterior column through a posterior approach to augment posterior instrumentation have been reported in the literature. We hypothesized that if anterior column support is an important adjunct to posterior short-segment instrumentation, then avoidance of axial load until sufficient anterior column healing occurs, allowing load-sharing with the implant, would improve spinal alignment at follow-up.

Methods: We conducted a retrospective cohort study in which consecutive patients who had instrumentation and fusion with the AO spinal fixator were immediately ambulated after surgery or had 4 weeks of bedrest. We measured kyphosis and wedge angles preoperatively, immediately postoperatively and at the time of final follow-up. We used radiologic measures to assess instrumentation and bone failure.

Results: We found significant differences in the mean loss of wedge and kyphosis angle correction between patients immediately ambulated and those who had 4 weeks of bedrest (0.71° v. -4.73° for wedge and 1.81° v. -6.55° for kyphosis, respectively). There was significant correlation between instrumentation and bone failure in both the immediate ambulation and bedrest groups.

Conclusion: Bedrest improves the maintenance of intraoperative sagittal alignment correction, which is in agreement with the theory that inadequate support of the anterior spinal column is the mechanism for loss of reduction when using short-segment instrumentation to treat burst fractures. Therefore, addressing the anterior column directly through anterior surgery or by employing novel techniques in posterior surgery is recommended if one of the goals of treatment is to maintain the sagittal correction achieved at the time of surgery. Trying to achieve this goal by addressing posterior implant design or bone quality alone will not be successful because instrumentation and bone failure occur together.

Contexte : La littérature fait état de réductions incomplètes attribuables au soutien insuffisant de la colonne antérieure lors de la pose de segments de plaques courts dans le traitement de la fracture-éclatement, et signale de nouvelles méthodes de soutien de la colonne antérieure par un abord postérieur afin d'augmenter l'ostéosynthèse par plaques par la voie postérieure. Nous avons émis l'hypothèse selon laquelle si le soutien de la colonne antérieure contribue substantiellement à l'ostéosynthèse obtenue au moyen de plaques postérieures courtes, l'évitement de toute mise en charge axiale jusqu'à l'atteinte d'un degré suffisant de consolidation de la colonne antérieure pour permettre la répartition de la charge sur l'implant améliorerait l'alignement spinal au moment du suivi.

Méthodes : Nous avons réalisé une étude de cohorte rétrospective au cours de laquelle des patients consécutifs ayant subi un montage par plaques et ostéosynthèse au moyen d'un fixateur spinal AO ont été mobilisés immédiatement après la chirurgie ou ont plutôt été confinés au lit pendant 4 semaines. Nous avons mesuré la cyphose et les angles de cunéiformisation en préopératoire, en post-opératoire immédiat et au moment du suivi final. Nous avons utilisé des mesures radiologiques pour évaluer l'échec au niveau du montage et de l'os.

Résultats : Nous avons découvert des différences significatives quant à la perte moyenne de cunéiformisation et la correction de l'angle de la cyphose selon que les patients avaient été immédiatement mobilisés ou étaient restés alités pendant 4 semaines (0,71 ° c. -4,73 ° pour la cunéiformisation et 1,81 ° c. -6,55 ° pour la cyphose, respectivement). On a noté une corrélation significative entre l'échec au niveau du montage et au

niveau osseux selon que les patients appartenaient au groupe immédiatement mobilisé et maintenu alité.

Conclusion : Le repos au lit améliore le maintien de la correction de l'alignement sagittal peropératoire, ce qui concorde avec la théorie selon laquelle un soutien inadéquat de la colonne spinale antérieure serait le mécanisme par lequel se produirait la réduction incomplète lors de l'utilisation de montages par fragments courts pour traiter les fractures-éclatements. Par conséquent, on recommande de traiter la colonne antérieure directement par l'abord chirurgical antérieur ou au moyen de nouvelles techniques utilisant l'abord postérieur si l'un des objectifs du traitement est de maintenir la correction sagittale obtenue au moment de la chirurgie. Tenter d'atteindre cet objectif en misant sur un implant conçu pour un abord postérieur ou sur la qualité osseuse seulement ne produira pas les résultats escomptés, parce que l'échec survient simultanément au niveau du montage et de l'os.

Burst fractures comprise about 15% of all thoracolumbar spinal injuries.¹⁻⁴ Treatment options include nonoperative management, posterior surgery, anterior surgery and combined anterior and posterior surgery, and there is much controversy over the optimal method of management.⁵

Nonoperative management, through a period of recumbency and/or the use of hyperextension casting or bracing, avoids the morbidity associated with surgical intervention, and its supporters cite acceptable functional results, particularly in patients with burst fractures who are neurologically intact on presentation.⁶⁻¹⁰ Alternatively, the advantages of surgery include a shorter period of bedrest and hospital admission, better correction of kyphotic deformity and the avoidance of late neurologic deterioration due to instability.^{2,3,11-13}

Earlier literature on surgical treatment supported the use of posterior instrumentation, citing excellent results with respect to spinal stability, postoperative neurologic improvement, anatomic alignment and low patient morbidity.¹⁴⁻¹⁷ Short-segment instrumentation such as the AO internal spinal fixator (Synthes) was considered to be more favourable than traditional Harrington rod posterior instrumentation and its modifications because short-segment instrumentation allowed stable fixation and reduction across all 3 columns and was limited only to adjacent spinal segments.¹⁴ Supporters of short-segment internal fixators have reported favourable results.^{1,4,18,19}

In more recent literature, detractors of a purely posterior approach have reported a loss of reduction due to inadequate support of the anterior spinal column.²⁰⁻²³ The pedicle screw-based implants have failed owing to loosening, bending or fracturing, resulting in a mean loss of kyphosis correction ranging from 9° to 11° in the reported pedicle screw studies.²⁴⁻²⁹ This has led to proponents for anterior strut grafting and instrumentation to treat burst fractures.³⁰⁻³³

Most recently, there have been reports of novel methods for support of the anterior column through a posterior approach to augment posterior short-segment instrumentation. Cho and colleagues³⁴ have reported on the use of polymethylmethacrylate (PMMA) vertebroplasty, Acosta

and colleagues³⁵ on the use of PMMA kyphoplasty, Verlaan and colleagues³⁶ on the use of calcium phosphate cement kyphoplasty and Li and colleagues³⁷ on the use of a titanium transpedicle body augments.

We sought to go back a step and demonstrate that the mechanism for loss of alignment correction in patients with burst fractures treated with short-segment posterior instrumentation was due to inadequate support of the anterior column. We hypothesized that if anterior column support is an important adjunct to posterior short-segment instrumentation in maintaining the sagittal correction achieved at the time of surgery, then avoidance of axial load until sufficient anterior column healing occurs, allowing load-sharing with the implant, would improve spinal alignment at follow-up. We tested this hypothesis by comparing maintenance of correction among patients with posterior instrumentation who were immediately ambulated after surgery to that among patients who had bedrest for the first 4 weeks after surgery to offload the spine.

METHODS

We retrospectively reviewed the medical records and radiographs of patients with burst fractures treated with posterior short-segment instrumentation and fusion by one of us (D.Y.) at our tertiary-care centre from October 1991 to June 2003. All patients underwent computed tomography (CT) scanning to confirm the diagnosis of a burst fracture as defined by Denis and colleagues.¹² Additional inclusion criteria were vertebral height loss greater than 50% or kyphosis greater than 20°, no neurologic deficit that would interfere with walking, no medical illnesses that would preclude operative intervention, no pre-existing diagnosis or active treatment for osteoporosis and sufficient pedicle size for posterior instrumentation. We obtained hospital ethics approval as a quality assurance review.

The treatment for all patients in both groups was posterior short-segment instrumentation with the AO spinal fixator and posterolateral fusion using autologous bone graft from the iliac crest, as previously described by Dick.¹⁶ In keeping with the early reports of maintaining deformity correction through posterior short-segment instrumentation

and fusion, we ambulated the initial cohort of patients immediately after surgery. Later, when reports of loss of correction began appearing in the literature, treatment changed to include an initial period of 4 weeks of bedrest to allow strengthening of the anterior column before axial loading. Patients in the bedrest group were allowed bathroom privileges and they were allowed to have the head of the bed raised up to 90°. They received deep vein thrombosis prophylaxis. All patients in both groups wore a spinal orthosis while ambulating for the first 3 months after their injuries.

We took radiographic measurements at 3 points during patient care: on admission, immediately after surgery and at final out-patient follow-up. We obtained the postoperative radiographs for patients in both cohorts using the same technique. We obtained the anteroposterior radiograph while patients were supine, and the lateral radiograph while the patients were in the lateral decubitus position. We measured kyphosis and wedge angles according to the method described by Denis and colleagues³⁸ (Fig. 1). Two observers measured all the radiographic parameters while blinded to the treatment groups, with differences settled by consensus.

There were no instances of broken screws. Therefore, to assess the contribution to the loss of kyphotic correction by implant bending (instrument failure) and screw migration (bone failure), we measured interscrew-to-rod ratios and screw-end plate-to-rod ratios (Fig. 2).

Statistical analysis

We entered the data into an Excel spreadsheet, and we used SPSS version 12.0.1 (SPSS Inc.) for our analysis. After descriptive analysis (frequencies, means, standard deviations),




Fig. 1. Kyphosis angle κ between a_1, b_1 and c_3, d_3 , reflecting the contribution of both bony collapse and disc damage to angular deformity. Wedge angle ω between a_2, b_2 and c_2, d_2 , reflecting solely the degree of bony collapse.

we assessed the data for normality of distribution using histograms and box plots. We used independent samples *t* tests to analyze the alignment data collected in the bedrest versus the ambulation groups. We used Pearson correlation to analyze the relation between bend and migration.

RESULTS

We reviewed the medical records and radiographs of 36 patients with burst fractures treated with posterior short-segment instrumentation, as confirmed on CT scans, from October 1991 to June 2003. Of these, 4 patients were lost to follow-up (3 owing to relocation and 1 owing to death from unrelated causes) before the final follow-up could be performed, which left 32 patients for inclusion in our study.

An initial 19 patients treated between October 1991 and September 1997 formed the immediate ambulation group, and 11 patients treated between October 1997 and June 2003 formed the bedrest group. Two patients, presenting in August 1998 and June 2001, respectively, requested immediate ambulation and thus were included in that group, bringing its number of participants to 21.

The immediate ambulation group comprised 16 men and 5 women with a mean age of 45 (range 17–76) years. There were 2 patients older than 70 years. Excluding these patients reduced the mean age of the group to 42 years. The mean follow-up was 18 (range 1–117) months. The location of the fractures were the T12 vertebra in 5 patients, L1 in 8, L2 in 6, L3 in 1 and L4 in 1. The mechanism of injury was motor vehicle collision in 7 patients, a fall from a height in 12, a direct blow from an object in 1 and no known trauma in 1.




Fig. 2. Interscrew-to-rod ratio, AB/c_3d_3 , as a measure of instrument failure. Screw-end plate-to-rod ratio, $(e + f)/c_3d_3$, as a measure of bone failure, where e and f are the distances from screw tips A and B to the superior and inferior end plates, respectively.

The bedrest group comprised 7 men and 4 women with a mean age of 42 (range 19–58) years. The mean follow-up was 16 (range 3–79) months. The location of the fractures were the T12 vertebra in 1 patient, L1 in 3, L2 in 4, L3 in 1 and L4 in 2. The mechanism of injury was motor vehicle collision in 6 patients, a fall from a height in 4 and a direct blow from a heavy object in 1.

Preoperative, postoperative and follow-up wedge angles and kyphosis angles were normally distributed, permitting the use of parametric statistics. The mean results for both groups are listed in Table 1. The change from the preoperative to the follow-up period was also normally distributed for both angles. There was no difference in the mean postoperative wedge angle between the immediate ambulation and bedrest groups (10.29° v. 11.27°, $t = -0.14$, $p = 0.66$). However, the mean change in wedge angle at final follow-up compared with preoperative angles was statistically significant at 0.71° in the immediate ambulation group and -4.73° in the bedrest group ($t = 2.56$, $p = 0.016$).

The difference in the mean postoperative kyphosis angle between the immediate ambulation and bedrest groups approached significance at 4.24° and -3.18°, respectively ($t = 1.75$, $p = 0.09$). However, the mean change in kyphosis angle at final follow-up compared with preoperative angles was 1.81° in the immediate ambulation group and -6.55° in the bedrest group ($t = 2.59$, $p = 0.015$).

We recognized that the follow-up was short in 3 patients (1 month in 2 patients and 2 months in another), therefore, we repeated our analysis excluding the data on these patients. Significance levels were similar despite the loss of power associated with the reduction in sample size: significance levels changed to $p = 0.017$ from $p = 0.016$ for the wedge angle and from $p = 0.015$ to $p = 0.012$ for the kyphosis angle, remaining statistically significant for both values.

Migration and bend were normally distributed, allowing the use of the parametric Pearson correlation to assess the association between the 2 measures. They were significant

and positively correlated at $r = 0.498$ ($p = 0.005$).

There were no decubitus ulcers, thromboemboli or other complications related to bedrest.

DISCUSSION

There were no differences in the mean postoperative wedge and kyphosis angles between the immediate ambulation and bedrest groups. However, in the former group, there was a failure to maintain the correction of sagittal alignment achieved intraoperatively, and there was a further loss of 0.71° of wedge angle and 1.81° of kyphosis angle at follow-up compared with the preoperative measurements. This is consistent with reports in the literature indicating that posterior short-segment instrumentation alone is insufficient in maintaining anatomic correction, with loss of kyphotic correction by the loosening, bending and fracturing of pedicle screw-based implants.^{20,23,30,32}

Sasso and colleagues¹⁷ noted that in patients who experience substantial compression of the anterior vertebral body, the disrupted anterior column lacks structural integrity and places significant bending loads on the implant. We hypothesized that avoidance of axial load until sufficient vertebral body healing occurs, providing anterior column support and allowing load-sharing with posterior short-segment instrumentation, would improve the maintenance of the intraoperative reduction. The significant difference that we observed between the ability of the immediate ambulation and bedrest groups to maintain the sagittal alignment correction obtained intraoperatively, with the bedrest cohort having a wedge angle that was 4.73° better at follow-up than preoperatively, supports the findings of Sasso and colleagues and our hypothesis.

Loss of reduction due to disc collapse has been reported by Lindsey and Dick²⁶ and Speth and colleagues;²⁷ the loss of reduction occurs because the intervertebral disc material is redistributed into the fractured vertebral body.^{39,40} This is important because it can contribute to the local kyphosis. As a result, we measured the kyphosis angle, which includes the vertebral body and the adjacent discs. There was a significant difference between the bedrest and immediate ambulation groups, with the former having a kyphosis angle that was 6.55° better at follow-up than preoperatively. We hypothesize that the period of bedrest allowed sufficient healing of the vertebral body to prevent redistribution of disc material with subsequent disc collapse, accounting for our finding.

We found that in the immediate ambulation and bedrest groups, there was loss of correction due to screw migration and bending of the instrumentation. Analysis demonstrated a positive correlation between migration and bend. This means that in patients with loss of correction, migration and bend occurred together. Therefore, the loss of reduction in posterior short-segment instrumentation is not due to implant or bone quality in isolation. This

Table 1. Preoperative, immediately postoperative and follow-up measurements of 32 patients with burst fractures, by treatment group

Time; measurement	Group; mean	
	Bedrest	Immediate ambulation
Preoperative		
Wedge angle	20.36°	14.57°
Kyphosis angle	8.91°	10.05°
Postoperative		
Wedge angle	11.27°	10.29°
Kyphosis angle	-3.18°	4.24°
Follow-up		
Wedge angle	15.64°	15.29°
Kyphosis angle	2.36°	11.86°

explains the finding of Alvine and colleagues²² that, despite progressive improvement in flexion bending endurance of the pedicle screws, there was still loss of sagittal alignment in follow-up when using variable screw plate instrumentation to treat burst fractures.

There were no complications due to bedrest, but we recognize the advantages of early ambulation and shortened hospital admission as well as our small sample size. Therefore, we present this study not to promote bedrest, but rather to demonstrate that the loss of alignment correction in patients with burst fractures treated with short-segment posterior instrumentation is due to inadequate support of the anterior column. Addressing inadequate support through novel techniques³⁴⁻³⁷ is an important adjunct to posterior implants in maintaining the sagittal correction achieved at the time of surgery.

Our study has limitations. Acquiring follow-up data on patients in a transitional population is difficult, and 4 of 36 patients (11%) were lost to follow-up. In addition, the change in practice over time to produce the 2 cohorts introduced a potential learning curve bias. This may have caused better results in the bedrest group, which was treated later. However, during this later time period, 2 patients requested immediate ambulation. One of these patients had better wedge (by 1°) and kyphosis (by 7°) angles at follow-up compared with preoperative measures. The other had worse wedge (by 6°) and kyphosis (by 5°) angles at follow-up compared with preoperative measures.

Our aim was to study the importance of the anterior column to the mechanism of failure of short segment instrumentation, not patient satisfaction or function with the use of bedrest. Therefore, we gathered radiologic parameters rather than long-term patient questionnaire data. Although our follow-up was relatively short, we believe our results are valid because radiographic analysis would define early on whether avoidance of axial load until sufficient vertebral healing occurred would improve sagittal alignment at follow-up.

CONCLUSION

Bedrest improves the maintenance of intraoperative sagittal alignment correction, consistent with the theory that inadequate support of the anterior spinal column is the mechanism for loss of reduction when using posterior short-segment instrumentation to treat burst fractures. Addressing the anterior column directly through anterior surgery or novel techniques in posterior surgery is recommended if one of the goals of treatment is to maintain the sagittal correction achieved at the time of surgery. Trying to achieve this goal by addressing posterior implant design or bone quality alone will not be successful because instrumentation and bone failure occur together.

Competing interests: None declared.

Contributors: Drs. Dang and Yen designed the study. Dr. Yen acquired the data, which Dr. Dang and Ms. Hopman analyzed. All authors wrote and reviewed the article and gave final approval for its publication.

References

1. Danisa OA, Shaffrey CI, Jane JA, et al. Surgical approaches for the correction of unstable thoracolumbar burst fractures: a retrospective analysis of treatment outcomes. *J Neurosurg* 1995;83:977-83.
2. Esses SI, Botsford DJ, Kostuik JP. Evaluation of surgical treatment for burst fractures. *Spine* 1990;15:667-73.
3. Denis F. The three column spine and its significance in the classification of acute thoracolumbar spinal injuries. *Spine* 1983;8:817-31.
4. Muller U, Berlemann U, Sledge J, et al. Treatment of thoracolumbar burst fractures without neurological deficit by indirect reduction and posterior instrumentation: bisegmental stabilization with monosegmental fusion. *Eur Spine J* 1999;8:284-9.
5. Verlaan JJ, Diekerhof CH, Buskens E, et al. Surgical treatment of traumatic fractures of the thoracic and lumbar spine: a systematic review of the literature on techniques, complications, and outcome. *Spine* 2004;29:803-14.
6. Cantor JB, Lebowitz NH, Garvey T, et al. Nonoperative management of stable thoracolumbar burst fractures with early ambulation and bracing. *Spine* 1993;18:971-6.
7. McAfee PC, Yuan HA, Frederickson BE, et al. The value of computed tomography in thoracolumbar fractures. *J Bone Joint Surg Am* 1983;65:461-73.
8. Mumford J, Weinstein JN, Spratt KF, et al. Thoracolumbar burst fractures. The clinical efficacy and outcome of nonoperative management. *Spine* 1993;18:955-70.
9. Weinstein JN, Collalto P, Lehmann TR. Thoracolumbar "burst" fractures treated conservatively: a long-term follow-up. *Spine* 1988;13:33-8.
10. Agus H, Kayali C, Arslantas M. Nonoperative treatment of burst-type thoracolumbar vertebral fractures: clinical and radiologic results of 29 patients. *Eur Spine J* 2005;14:536-40.
11. Chow GH, Nelson BJ, Gebhard JS, et al. Functional outcome of thoracolumbar burst fractures managed with hyperextension casting or bracing and early mobilization. *Spine* 1996;21:2170-5.
12. Denis F, Armstrong GW, Searis K, et al. Acute thoracolumbar burst fractures in the absence of neurologic deficit. A comparison between operative and nonoperative treatment. *Clin Orthop Relat Res* 1984;189:142-9.
13. Noel SH, Keene JS, Rice WL. Improved postoperative course after spinous process segmental instrumentation of thoracolumbar fractures. *Spine* 1991;16:132-6.
14. Akalm S, Kis M, Benli IT, et al. Results of the AO spinal internal fixator in the surgical treatment of thoracolumbar burst fractures. *Eur Spine J* 1994;3:102-6.
15. Aebi M, Etter C, Kehl T, et al. Stabilization of the lower thoracic and lumbar spine with the internal spinal skeletal fixation system. Indications, techniques, and first results of treatment. *Spine* 1987;12:544-51.
16. Dick W. The "fixateur interne" as a versatile implant for spine surgery. *Spine* 1987;12:882-900.
17. Sasso RC, Cotler HB, Reuben JD. Posterior fixation of thoracic and lumbar spine fractures using DC plates and pedicle screws. *Spine* 1991;16:S134-9.
18. Andreychik DA, Alander DH, Senica KM, et al. Burst fractures of the second through fifth lumbar vertebrae. Clinical and radiographic results. *J Bone Joint Surg Am* 1996;78:1156-66.

19. Zindrick MR, Lorenz MA. The use of intrapedicular fixation systems in the treatment of thoracolumbar and lumbosacral fractures. *Orthopedics* 1992;15:337-41.
20. Razak M, Mahmud MM, Hyzan MY, et al. Short segment posterior instrumentation, reduction and fusion of unstable thoracolumbar burst fractures — a review of 26 cases. *Med J Malaysia* 2000;55:9-13.
21. Parker JW, Lane JR, Karaikovic EE, et al. Successful short-segment instrumentation and fusion for thoracolumbar spine fractures. *Spine* 2000;25:1157-69.
22. Alvine GF, Swain JM, Asher MA, et al. Treatment of thoracolumbar burst fractures with variable screw placement or isola instrumentation and arthrodesis: case series and literature review. *J Spinal Disord Tech* 2004;17:251-64.
23. Wood K, Butterman G, Mehdod A, et al. Operative compared with nonoperative treatment of a thoracolumbar burst fracture without neurological deficit. *J Bone Joint Surg Am* 2003;85:773-81.
24. Gurwitz GS, Dawson JM, McNamara MJ, et al. Biomechanical analysis of three surgical approaches for lumbar burst fractures using short-segment instrumentation. *Spine* 1993;18:977-82.
25. McLain RF, Sparling E, Benson DR. Early failure of short-segment pedicle instrumentation for thoracolumbar fractures. *J Bone Joint Surg Am* 1993;75:162-7.
26. Lindsey RW, Dick W. The fixateur interne in the reduction and stabilization of thoracolumbar spine fractures in patients with neurologic deficit. *Spine* 1991;16:S140-5.
27. Speth MJ, Oner FC, Kadic MA, et al. Recurrent kyphosis after posterior stabilization of thoracolumbar fractures. *Acta Orthop Scand* 1995;66:406-10.
28. Shen WJ, Liu TJ, Shen YS. Nonoperative treatment versus posterior fixation for thoracolumbar junction burst fractures without neurologic deficit. *Spine* 2001;26:1038-45.
29. Alanay A, Acaroglu E, Yazici M, et al. Short-segment pedicle instrumentation of thoracolumbar burst fractures. Does transpedicular intracorporeal grafting prevent early failure? *Spine* 2001;26:213-7.
30. Kaneda K, Taneichi H, Abumi K, et al. Anterior decompression and stabilization with the Kaneda device for thoracolumbar burst fractures associated with neurological deficit. *J Bone Joint Surg Am* 1997;79:69-83.
31. McDonough PW, Davis R, Tribus C, et al. The management of acute thoracolumbar burst fractures with anterior corpectomy and Z-Plate fixation. *Spine* 2004;29:1901-8.
32. Wood KB, Bohn D, Mehdod A. Anterior versus posterior treatment of stable thoracolumbar burst fractures without neurologic deficit: a prospective, randomized study. *J Spinal Disord Tech* 2005;18(Suppl 1):S15-23.
33. McCormack T, Karaikovic E, Gaines RW. The load sharing classification of spine fractures. *Spine* 1994;19:1741-4.
34. Cho DY, Lee WY, Sheu PC. Treatment of thoracolumbar burst fractures with polymethyl methacrylate vertebroplasty and short-segment pedicle screw fixation. *Neurosurgery* 2003;53:1354-61.
35. Acosta FL, Aryan HE, Taylor WR, et al. Kyphoplasty-augmented short-segment pedicle screw fixation of traumatic thoracolumbar burst fractures: initial clinical experience and literature review. *Neurosurg Focus* 2005;18:E9.
36. Verlaan JJ, Dhert WJA, Verbout AJ, et al. Balloon vertebroplasty in combination with pedicle screw instrumentation: a novel technique to treat thoracic and lumbar burst fractures. *Spine* 2005;30:E73-9.
37. Li KC, Hsieh C-H, Lee C-Y, et al. Transpedicle body augments: a further step in treating burst fractures. *Clin Orthop Relat Res* 2005;436:119-25.
38. Denis F, Ruiz H, Searls K. Comparison between square-ended distraction rods and standard round-ended distraction rods in the treatment of thoracolumbar spinal injuries — a statistical analysis. *Clin Orthop Relat Res* 1984;189:162-7.
39. Oner FC, van der Rijt RR, Ramos LM, et al. Changes in the disc space after fractures of the thoracolumbar spine. *J Bone Joint Surg Br* 1998;80:833-9.
40. Verlaan JJ, van Helden WH, Oner FC, et al. Balloon vertebroplasty with calcium phosphate cement augmentation for direct restoration of traumatic thoracolumbar vertebral fractures. *Spine* 2002;27:543-8.



Canadian Journal
of **Surgery**

Online manuscript
submission and peer review

NOW AVAILABLE FOR CJS at
<http://mc.manuscriptcentral.com/cjs>