

A day in the life of emergency general surgery in Canada: a multicentre observational study

Kristin DeGirolamo, MD
 Karan D'Souza, BSc
 Sameer Apte, MD
 Chad G. Ball, MD
 Christopher Armstrong, MD
 Artan Reso, MD
 Sandy Widder, MD
 Sarah Mueller, MD
 Lawrence M. Gillman, MD
 Ravinder Singh, MD
 Rahima Nenshi, MD
 Kosar Khwaja, MD
 Samuel Minor, MD
 Chris de Gara, MB, MS
 S. Morad Hameed, MD

Accepted Nov. 17, 2017; Published online
 June 1, 2018

Correspondence to:

M. Hameed
 Trauma Services, Vancouver General
 Hospital
 855 West 12th Ave
 Vancouver BC V5Z 1M9
 morad.hameed@vch.ca

DOI: 10.1503/cjs.013517

Background: Emergency general surgery (EGS) services are gaining popularity in Canada as systems-based approaches to surgical emergencies. Despite the high volume, acuity and complexity of the patient populations served by EGS services, little has been reported about the services' structure, processes, case-mix or outcomes. This study begins a national surveillance effort to define and advance surgical quality in an important and diverse surgical population.

Methods: A national cross-sectional study of EGS services was conducted during a 24-hour period in January 2017 at 14 hospitals across 7 Canadian provinces recruited through the Canadian Association of General Surgeons Acute Care Committee. Patients admitted to the EGS service, new consultations and off-service patients being followed by the EGS service during the study period were included. Patient demographic information and data on operations, procedures and complications were collected.

Results: Twelve sites reported resident coverage. Most services did not include trauma. Ten sites had protected operating room time. Overall, 393 patient encounters occurred during the study period (195/386 [50.5%] operative and 191/386 [49.5%] nonoperative), with a mean of 3.8 operations per service. The patient population was complex, with 136 patients (34.6%) having more than 3 comorbidities. There was a wide case-mix, including gallbladder disease (69 cases [17.8%]) and appendiceal disease (31 [8.0%]) as well as complex emergencies, such as obstruction (56 [14.5%]) and perforation (23 [5.9%]).

Conclusion: The characteristics and case-mix of these Canadian EGS services are heterogeneous, but all services are busy and provide comprehensive operative and nonoperative care to acutely ill patients with high levels of comorbidity.

Contexte : Les services de chirurgie générale d'urgence (CGU) gagnent en popularité au Canada en tant qu'approches systémiques aux urgences chirurgicales. Malgré le volume élevé, le caractère urgent et la complexité des populations de patients desservies en CGU, peu de rapports ont porté sur la structure, les processus, les clientèles ou les résultats de ces services. La présente étude instaure une démarche de surveillance nationale qui servira à définir et à améliorer la qualité des chirurgies destinées à cette population importante et hétérogène.

Méthodes : Une étude transversale nationale sur les services de CGU a été réalisée sur une période de 24 heures en janvier 2017 dans 14 hôpitaux de 7 provinces canadiennes recrutés par l'entremise du comité pour les soins aigus de l'Association canadienne des chirurgiens généraux. On y a inclus les patients admis dans les services de CGU, les nouvelles consultations et les patients de l'extérieur suivis par les services de CGU pendant la période de l'étude. On a recueilli les caractéristiques démographiques des patients et les données sur les interventions, les procédures et les complications.

Résultats : Douze sites ont fait état de la couverture assurée par les résidents. La plupart des services ont exclu la traumatologie. Dix sites disposaient de temps protégé au bloc opératoire. En tout, 393 rencontres avec des patients ont eu lieu pendant la période de l'étude (195/386 [50,4 %] chirurgicales, 191/386 [49,5 %] non chirurgicales), avec une moyenne de 3,8 chirurgies par service. La population regroupait des cas complexes : 136 patients (34,6 %) présentaient plus de 3 comorbidités. La clientèle était diversifiée et comprenait des cas de maladie de la vésicule biliaire (69 cas [17,8 %]) et de l'appendice (31 [8,0 %]), de même que des situations d'urgence délicates, telle qu'obstruction (56 [14,5 %]) et perforation (23 [5,9 %]).

Conclusion : Leurs caractéristiques et leurs clientèles sont hétérogènes, mais les services de CGU sont tous achalandés et ils offrent tous des soins chirurgicaux et non chirurgicaux complets à des patients gravement malades porteurs d'importantes comorbidités.

General surgical emergencies are common. They may account for 25%–50% of a general surgeon’s overall workload and 7% of all hospital admissions.^{1,2} In 2006, the Institute of Medicine declared non-trauma surgical emergencies as one of the main stressors on emergency departments in the United States.³ Patients facing surgical emergencies are often critically ill with numerous comorbidities. Timely, comprehensive intervention can prevent devastating long-term complications.³ Traditionally, on-call surgeons have been required to balance emergency surgical care with a busy elective subspecialty service, including rounds with elective patients, visiting patients within clinic and daytime subspecialty operating.

In Canada, there has been a trend toward consolidating emergency general surgery (EGS) onto dedicated services.^{1,2} Emergency general surgery services have been implemented with the intent of providing a dedicated hospital-based service specifically for the care of general surgical emergencies.^{4,5} This allows comprehensive care to be provided while also creating stronger platforms for quality improvement and optimization of resource use. There are limited data on the overall case-mix of these novel EGS services and their varied structures, processes and out-

comes.^{6–8} The emergence of EGS systems represents an opportunity to understand and improve processes of care that serve this complex, resource-intensive surgical population. More research is required on the true acuity, complexity and diversity of EGS services and systems, as well as on the roles these services play in supporting surgical rescue within acute care health systems.

This study captures a snapshot at a single point in time of both the case-mix and workflow during a typical day in EGS across Canada. We hypothesized that Canadian EGS services face highly acute, complex, diverse case-mixes and that service models have evolved in unique ways in response to specific requirements within their local environments. Shared insights about case-mix and service delivery will inform the next generation of developments in quality improvement and health system design.

METHODS

Design

The Canadian Association of General Surgeons formed the Acute Care Committee from a group of surgeons providing trauma and EGS care across Canada. This inclusive




Fig. 1. Participating sites.

committee outlined a road map for multicentre EGS research, starting with an environmental scan of EGS services across the country. Such a study creates a strong foundation for future clinical and systems studies in EGS, characterizes future participating sites, identifies opportunities for collaboration, helps to disseminate best practices and serves as a gap analysis to identify new research directions. Many members of the committee are EGS site leaders, which was an advantage for study design, institutional review and data acquisition. The protocol for this cross-sectional study was approved by the ethics review board at the study lead site (Vancouver General Hospital) and subsequently at each of the participating hospitals.

Service structure

A survey of EGS service leaders was used to characterize the structure of different services at each of 14 hospitals across 7 provinces in Canada (Fig. 1). Site leads were asked to complete a database sheet to describe key features of their EGS services. Examples of structural features include the use of a “surgeon of the week” model, the presence of a dedicated EGS service, the availability of protected operating room time for EGS cases and the ability to use EGS operating room time for elective cases. Site leads were also asked whether all general and subspecialty surgeons participated and about the involvement of resident coverage, the inclusion of trauma patients and the existence of a formalized handover process between surgeons at shift change.

Case-mix

Patients assessed by EGS services at participating sites on Jan. 10, 2017, formed the study cohort, and the EGS teams on call from 7 am on Jan. 10, 2017, to 7 am on Jan. 11, 2017, were briefed about the protocol. Patients currently admitted to the EGS service, new consultations and off-service patients (admitted to the hospital but not under the direct care of an EGS service) being followed by the EGS service during the 24-hour study period were included in the study. Trauma patients and patients discharged from the EGS service before 7 am on Jan. 10, 2017, were excluded.

Participating sites collected data using standardized case report forms and a summary sheet (Appendix 1, available at canjsurg.ca/013517-a1). Data were collected by study personnel at each site under the supervision of the site lead (coauthors of this paper), and the primary author or the site lead entered the data. Patient demographic information such as age, sex, comorbidities and previous operations was collected, as were all operative and procedural data, and data on intraoperative and postoperative complications. Data were verified with the site leads individually, and all perceived discrepancies and questions were resolved in a second round of communication with all sites.

Data analysis

Site leads tabulated data from the case report forms and summary sheets, and these were merged in Microsoft Excel version 15.18. Subsequently, basic statistics were conducted and condensed into tables.

RESULTS

Service structure

Emergency general surgery services across Canada reported using different structural models for the delivery of care (Table 1). All participating sites had a dedicated EGS service with a “surgeon of the week” model, whereby 1 surgeon suspends his or her elective practice for an entire week to focus on EGS patients during the day, sharing overnight call responsibilities. Ten sites (71%) had protected operating room time, and 6 (43%) were permitted to use protected time for elective cases. In contrast to the situation in the US, where trauma and EGS are often combined in single services,^{9,10} only 4 Canadian sites (29%) included trauma patients under the scope of EGS services. Twelve sites (86%) reported resident coverage, and all had a formal handover process.

Case-mix

A total of 393 patients were assessed across the 14 sites during the study period. The EGS service was newly consulted to assess 112 patients, of whom 76 (67.8%) were admitted by the EGS service within the 24-hour period (Table 2). Of patients previously admitted to the EGS service, 40 (13.0%) were cared for in an intensive care unit setting, and 10 (3.2%) had open abdomens. Emergency general surgery teams operated in 53 cases during the study period, 28 (53%) of which were completed laparoscopically. When considering all patients on the EGS census, 195 cases (50.5%) were operative (Table 3). Of the operative cases, 88 (44.9%) were laparoscopic, with a conversion rate of 5.7%. There was 1 intraoperative complication across all sites. A total of 109 (57.1%) of the nonoperative cases were managed with antibiotic regimens.

The mean patient age was 59.1 (standard deviation 4.9) years, and 186 (47.3%) were female. Most patients assessed by the EGS service had complex conditions: 207 (52.7%) had 1–3 comorbidities, and 136 (34.6%) had more than 3 comorbid conditions (Table 4). In addition, there was a wide range of presenting problems and final diagnoses evaluated and managed by EGS teams (Table 5). The most common diagnoses were gallbladder disease (69 patients [17.8%]), gastric/intestinal obstruction (56 [14.5%]) and appendiceal disease (31 [8.0%]). The diagnoses that most increased the average length of stay in hospital were neoplasms, pancreatic disease and intestinal obstruction.

DISCUSSION

This study represents a unique snapshot of the complex care that is provided on an average day at EGS services across 14 Canadian hospitals. Consequently, it provides a novel perspective of a large, complex, resource-intensive patient population, for which limited reporting exists within the literature.^{1,9}

The development of EGS services in Canada emerged organically as a way to address the growing complexity of EGS. Implementing these services has highlighted aims to improve access to care, patient outcomes and safety, while enhancing efficiency of service delivery, surgeon satisfaction and educational opportunities.¹¹ As shown by the variation in models adopted at the 14 participating sites, hospitals have created services structured to meet these

Table 1. Description of emergency general surgery service structures and processes across Canada

Hospital service	Dedicated service	“Surgeon of the week” model	Protected operating room time	Resident coverage	Trauma included	All surgeons participate	Formal handover	Elective cases in protected time
Vancouver General Hospital	X	X	X	X		X	X	X
Foothills Medical Centre, Calgary	X	X		X		X	X	
South Health Campus, Calgary	X	X	X			X	X	
Rockyview General Hospital, Calgary	X	X	X	X		X	X	
University of Alberta Hospital, Edmonton	X	X	X	X		X	X	X
Royal University Hospital, Saskatoon	X	X	X	X	X	X	X	X
St. Paul’s Hospital, Saskatoon	X	X	X	X		X	X	X
St. Boniface Hospital, Winnipeg	X	X	X	X		X	X	
Health Sciences Centre, Winnipeg	X	X		X	X	X	X	
St. Joseph’s Healthcare, Hamilton, Ont.	X	X		X		X	X	
North Bay Regional Health Centre, North Bay, Ont.	X	X			X		X	
Royal Victoria Hospital, Montréal	X	X		X		X	X	
Montreal General Hospital	X	X	X	X		X	X	X
Queen Elizabeth II Health Science Centre, Halifax	X	X	X	X	X	X	X	X

Table 2. Cohorts of emergency general surgery patients, by hospital service

Hospital service	No. (%) of patients					
	Total followed by EGS service	Seen in consultation	Admitted	Admitted to intensive care unit	Open abdomen	Following without admission
Vancouver General Hospital	33	9	7	4 (12)	1 (3)	4 (12)
Foothills Medical Centre	36	15	11	3 (8)	1 (3)	1 (3)
South Health Campus	24	7	5	3 (12)	0 (0)	3 (12)
Rockyview General Hospital	23	12	9	0 (0)	0 (0)	7 (30)
University of Alberta Hospital	10	4	3	6 (60)	2 (20)	5 (50)
Royal University Hospital	10	6	3	0 (0)	3 (30)	1 (10)
St. Paul’s Hospital	19	10	6	1 (5)	0 (0)	5 (26)
St. Boniface Hospital	35	7	7	0 (0)	0 (0)	0 (0)
Health Sciences Centre	28	10	8	10 (36)	1 (4)	3 (11)
St. Joseph’s Healthcare	28	4	2	3 (11)	0 (0)	10 (36)
North Bay Regional Health Centre	13	8	6	4 (31)	0 (0)	5 (38)
Royal Victoria Hospital	12	2	1	3 (25)	0 (0)	0 (0)
Montreal General Hospital	7	4	2	0 (0)	0 (0)	1 (14)
Queen Elizabeth II Health Science Centre	30	14	6	3 (10)	2 (7)	7 (23)
Total	308	112	76	40 (13.0)	10 (3.2)	52 (16.9)

EGS = emergency general surgery.

Table 3. Summary of operative and nonoperative management of emergency general surgery patients by hospital service*

Hospital service	No. of laparoscopic procedures	No. of open procedures	Total no. (%) of procedures	No. of laparoscopic to open procedure conversions	Total no. (%) nonoperative	No. of patients treated nonoperatively who received antibiotics
Vancouver General Hospital	8	17	25 (52)	1	23 (48)	8
Foothills Medical Centre	10	9	19 (56)	0	15 (44)	13
South Health Campus	8	6	14 (52)	0	13 (48)	10
Rockyview General Hospital	13	3	16 (67)	1	8 (33)	4
University of Alberta Hospital	2	13	15 (60)	0	10 (40)	5
Royal University Hospital	4	3	7 (41)	0	10 (59)	9
St. Paul's Hospital	6	12	18 (69)	1	8 (31)	5
St. Boniface Hospital	12	7	19 (54)	1	16 (46)	11
Health Sciences Centre	4	12	16 (61)	0	11 (39)	6
St. Joseph's Healthcare	9	4	13 (45)	1	16 (55)	10
North Bay Regional Health Centre	1	6	7 (39)	0	11 (61)	7
Royal Victoria Hospital	4	2	6 (30)	0	14 (70)	8
Montreal General Hospital	0	1	1 (8)	0	12 (92)	4
Queen Elizabeth II Health Science Centre	7	12	19 (44)	0	24 (56)	9
Total	88	107	195 (50.5)	5	191 (49.5)	109

*For 7 patients, no definitive management plan (neither operative nor nonoperative) was assigned during the study period.

Table 4. Age, sex and comorbidity status of emergency general surgery patients, by hospital service

Hospital service	No. of acute patients	Mean age, yr	Female sex, no. (%)	> 3 comorbidities, no. (%)
Vancouver General Hospital	48	62.5	27 (56)	17 (35)
Foothills Medical Centre	36	58.2	15 (42)	17 (47)
South Health Campus	27	56.6	11 (41)	3 (11)
Rockyview General Hospital	28	59.8	14 (50)	2 (7)
University of Alberta Hospital	25	63.5	8 (32)	18 (72)
Royal University Hospital	17	50.1	12 (70)	3 (18)
St. Paul's Hospital	26	59.1	14 (54)	6 (23)
St. Boniface Hospital	35	64.1	12 (34)	16 (46)
Health Sciences Centre	28	51.5	16 (57)	9 (32)
St. Joseph's Healthcare	29	68.1	10 (34)	21 (72)
North Bay Regional Health Centre	18	58.5	7 (39)	9 (50)
Royal Victoria Hospital	20	57.5	8 (40)	4 (20)
Montreal General Hospital	13	62.4	8 (62)	3 (23)
Queen Elizabeth II Health Science Centre	43	56.1	25 (58)	8 (19)

Table 5. Case-mix and resource use as reflected by duration of hospital stay at the time of the study

Final diagnosis	No. (%) of cases	Days since admission	
		Mean \pm SD	Median (range)
Gallbladder disease	69 (17.6)	5.3 \pm 8.0	4.0 (1–61)
Intestinal obstruction	56 (14.2)	11.9 \pm 22.1	4.0 (1–146)
Appendiceal disease	31 (7.9)	4.2 \pm 8.0	1.5 (1–40)
Neoplasm	25 (6.4)	21.7 \pm 41.5	7.0 (1–210)
Perforation	23 (5.8)	10.7 \pm 11.0	6.0 (1–44)
Diverticular disease	19 (4.8)	8.0 \pm 12.0	4.0 (1–50)
Gastrointestinal bleed	18 (4.6)	8.1 \pm 13.9	2.5 (1–53)
Hernia disease	16 (4.1)	4.2 \pm 4.0	3.0 (1–17)
Skin/soft-tissue infection	16 (4.1)	6.1 \pm 4.3	6.5 (1–14)
Pancreatic disease	15 (3.8)	15.4 \pm 24.6	5.0 (1–81)
Other*	105 (26.7)	—	—

SD = standard deviation.
*Includes abscess, anorectal disease, breast disease and sepsis.

objectives based on their local context. The number of surgeons within a department, diversity in the continuum of learners at a site, proximity of tertiary trauma care, and executive administration and financial support are a few potential factors that can affect the organization of an EGS service. The unpredictable pace of emergency surgery has made protected operating room time an important part of delivering timely, safe care. However, our findings suggest that there remains a gap in providing allocated time to emergency general surgeons. Furthermore, autonomy in deciding how unused resources are assigned is lacking.

Most studies on EGS have focused on processes and outcomes for appendicitis and cholecystitis.^{8,12-14} Although these conditions account for a substantial part of the activity of EGS services, they do not reflect the breadth, intensity or resource consumption of a modern EGS practice. In our national EGS sample, appendicitis and cholecystitis together accounted for only 26% of patients and only 10% of hospital stays. In contrast, cancer accounted for close to 23% of hospital days and, therefore, a considerable portion of EGS service activity.

Our findings confirm that Canadian EGS services are complex. EGS service team members not only provide operative care but also manage complicated nonoperative cases. Nationally, 49% of patients were managed nonoperatively, with the Montreal General Hospital site having the highest number of patients receiving nonoperative care. Although these patients often require several resource-intensive days in hospital, their care is not well documented in the EGS literature. The present study shows an opportunity for research that aims to understand and improve the processes and outcomes of care for EGS patients managed nonoperatively.

The complexity of EGS may, in part, be reflected by the extent of comorbidities and the frequent need for critical care. At 1 service, 72% of patients had more than 3 comorbid conditions. The fact that all of these patients had been considered for, or had undergone, major emergency operations is evidence that EGS services bear the responsibilities not only of assessment, diagnosis and resuscitation but also of preoperative optimization, operative intervention and postoperative care. This care is delivered to vulnerable patients with complex medical conditions over rapid time frames. Furthermore, the fact that 13.0% of patients in our study were cared for in the intensive care unit shows the critical status of many EGS patients. This intersection of aggressive surgical care and patients with complex medical conditions is a daily reality on modern EGS services. Consequently, these services provide an opportunity for multicentre quality improvement, guideline development and promotion of best practices. There is an urgent need for the next generation of EGS research to explore the intricacies of service delivery more fully.

The implications of a national EGS research network are far-reaching. Our study shows that EGS patient populations are large, complex and resource intensive, and that the structures and processes of EGS care are variable. A national EGS research network will be able to share experiences and define best practices and will serve as a forum to make these experiences and best practices more universal. The network has already defined a research and quality-improvement road map, with the next series of studies underway. Future studies will focus on processes of care, complex operative care, determinants of complications and death, benchmarks of quality and surgical education in EGS. It is hoped that ultimately this work will lead

to a national EGS database and research strategy dedicated to analyzing diversity in the Canadian EGS experience. With that national initiative, the structure, processes and outcomes of EGS service delivery can be optimized.

Strengths and limitations

This study was a one-time snapshot of 24 hours of care by EGS services at major Canadian hospitals. It was an observational project that did not capture every hospital with or without a formalized EGS service (mostly owing to difficulty in identifying these services and engaging them in a national study) and therefore could not provide a comprehensive view of Canadian EGS practice. However, despite its methodological limitations, the study is proof of the concept that it is possible to bring emerging EGS services and patient-level insights into a national research network. The fact that the study was completed rapidly shows that national collaboration on research protocol development, coordination of ethics review board applications, creation of data-sharing agreements, and shared data analysis, interpretation and reporting are highly feasible.

CONCLUSION

Canadian EGS services are at the centre of Canadian acute care. They are busy intake services for extremely vulnerable patients with a spectrum of complex, life-threatening conditions including abdominal sepsis, intestinal obstruction and cancer. Emergency general surgery services optimize perioperative and operative care along rapid timelines and often use nonoperative approaches as well. They act as rescue services, supporting patient care on other services, and frequently provide care in intensive care units. Ongoing national research collaborations will continue to shed light on the structures, processes and outcomes of these important new services and will identify new opportunities to improve patient care and system performance.

Acknowledgements: The authors thank the Canadian Association of General Surgeons Acute Care Committee for participating in this study and volunteering their hospital data and research staff to help make this project possible. The authors acknowledge Jillian Aquino for her design and creation of the figure.

Affiliations: From the Department of Surgery, Faculty of Medicine, University of British Columbia, Vancouver, BC (DeGirolamo, Hameed); the Faculty of Medicine, University of British Columbia, Vancouver, BC (D'Souza); the Department of Surgery, University of Alberta, Edmonton, Alta. (Apte, Widder, de Gara); the Department of Surgery, University of Calgary, Calgary, Alta. (Ball, Armstrong, Reso); the Department of Surgery, University of Saskatchewan, Saskatoon, Sask. (Mueller); the Department of Surgery, University of Manitoba, Winnipeg, Man. (Gillman); the Department of Surgery, Northern Ontario School of Medicine, North Bay, Ont. (Singh); the Department of Surgery, McMaster University, Hamilton, Ont. (Nenshi); the Department of Surgery, McGill University, Montréal, Que. (Khawaja); the Department of Surgery, Dalhousie University, Halifax, NS (Minor); and the Department of Trauma Services, Vancouver General Hospital, Vancouver, BC (Hameed).

Competing interests: None declared.

Contributors: K. DeGirolamo, K. D'Souza, S. Apte, C.G. Ball and S.M. Hameed designed the study. All authors acquired the data, which K. DeGirolamo, K. D'Souza, S. Apte, C.G. Ball and S.M. Hameed analyzed. K. DeGirolamo, K. D'Souza, S. Apte, C.G. Ball and S.M. Hameed wrote the article, which all authors reviewed and approved for publication.

References

1. Ball CG, Hameed SM, Brennenman FD. Acute care surgery: a new strategy for the general surgery patients left behind. *Can J Surg* 2010; 53:84-5.
2. Becher RD, Hoth JJ, Miller PR, et al. A critical assessment of outcomes in emergency versus nonemergency general surgery using the American College of Surgeons National Surgical Quality Improvement Program database. *Am Surg* 2011;77:951-9.
3. Santry HP, Pringle PL, Collins CE, et al. A qualitative analysis of acute care surgery in the United States: it's more than just "a competent surgeon with a sharp knife and a willing attitude." *Surgery* 2014;155:809-25.
4. Chana P, Burns EM, Arora S, et al. A systematic review of the impact of dedicated emergency surgical services on patient outcomes. *Ann Surg* 2016;263:20-7.
5. Nagaraja V, Eslick GD, Cox MR. The acute surgical unit model versus the traditional "on call" model: a systematic review and meta-analysis. *World J Surg* 2014;38:1381-7.
6. Faryniuk AM, Hochman DJ. Effect of an acute care surgical service on the timeliness of care. *Can J Surg* 2013;56:187-91.
7. Anantha RV, Parry N, Vogt K, et al. Implementation of an acute care emergency surgical service: a cost analysis from the surgeon's perspective. *Can J Surg* 2014;57:E9-14.
8. Wanis KN, Hunter AM, Harington MB, et al. Impact of an acute care surgery service on timeliness of care and surgeon satisfaction at a Canadian academic hospital: a retrospective study. *World J Emerg Surg* 2014;9:4.
9. Napolitano LM, Fulda GJ, Davis KA, et al. Challenging issues in surgical critical care, trauma, and acute care surgery: a report from the Critical Care Committee of the American Association for the Surgery of Trauma. *J Trauma Inj Infect Crit Care* 2010;69:1619-33.
10. Moore E. Acute care surgery: the safety net hospital model. *Surgery* 2007;141:297-8.
11. Warnock GL. Dynamic growth of the acute care surgery model [editorial]. *Can J Surg* 2010;53:76-7.
12. Qureshi A, Smith A, Wright F, et al. The impact of an acute care emergency surgical service on timely surgical decision-making and emergency department overcrowding. *J Am Coll Surg* 2011;213:284-93.
13. Krouchev R, Champagne-Parent G, Joos E, et al. The effect of an acute care surgery service on the management of appendicitis. *J Am Coll Surg* 2014;219:e153.
14. Lim DW, Ozegovic D, Khadaroo RG, et al. Impact of an acute care surgery model with a dedicated daytime operating room on outcomes and timeliness of care in patients with biliary tract disease. *World J Surg* 2013;37:2266-72.