

RESPONSE TO: PROPOFOL ADMINISTRATION BY ENDOSCOPISTS VERSUS ANESTHESIOLOGISTS IN GASTROINTESTINAL ENDOSCOPY: A SYSTEMATIC REVIEW AND META-ANALYSIS OF PATIENT SAFETY OUTCOMES

Daza and colleagues¹ performed a meta-analysis of the literature to compare the safety of the administration of propofol by nonanesthesiologists to low-risk patients (American Society of Anesthesiologists score of I or II) undergoing endoscopy procedures. The possible complications of sedation for endoscopy are well described in their article. However, their review did not differentiate between endoscopies done in hospitals and those done in out-of-hospital premises. The resources to manage complications of propofol in hospital may be significantly different than those in other facilities. Also, it is not clear from their review what standards of care were in place at the institutes where the reports included in the meta-analysis originated. The standards of care in place for the premises and the adherence to these standards by the providers could affect the data significantly.

Regulations are essential for patient safety and it is important for endoscopists to know what the requirements for sedation are and who else must be a member of the team. The Out-of-Hospital Premises Inspection Program of the College of Physicians and Surgeons of Ontario was initiated in 2010 and maintains standards for the safe delivery of sedation during endoscopic procedures.² These standards were developed through extensive consultation with physicians and surgeons who had broad experience and expertise in this work. The standards are consistently monitored and updated as the practice standards of optimum patient care evolve. Many other Canadian provinces and territories have similar

standards and regulation to ensure the safety and quality of patient care.

Owing to the potential for rapid and profound changes in anesthetic depth and the lack of antagonist medications, patients who receive propofol must receive care that is consistent with deep sedation even if moderate sedation is intended. These medications must be administered by a physician qualified to provide deep sedation. The physician administering propofol does not have to be an anesthesiologist, but does need to be qualified to administer deep sedation according to regulation and have the education and experience to manage the potential medical complications.

The standards² further specify that if the physician administering the sedation is also performing the procedure, the patient has to be attended by a second individual (physician, respiratory therapist, registered nurse or anesthesia assistant) who is not assisting in the procedure and who is trained to monitor patients undergoing sedation.

The critical issue for endoscopic procedures is not the administration of propofol by an anesthesiologist versus an endoscopist, but rather the capability of the physician administering propofol to manage its complications, the monitoring of the patient to detect complications and the resources to manage those complications.

Robert Byrick, MD; Dennis Pitt, MD, MEd

Affiliations: From the Department of Anesthesia, University of Toronto, Toronto, Ont. (Byrick); and the Department of Surgery, University of Ottawa, Ottawa, Ont. (Pitt).

Competing interests: None declared.

DOI: 10.1503/cjs.011618

References

1. Daza JF, Tan CM, Fielding RJ, et al. Propofol administration by endoscopists versus anesthesiologists in gastrointestinal endoscopy: a systemic review and meta-analysis of patient safety outcomes. *Can J Surg* 2018;61:226-36.

2. College of Physicians and Surgeons of Ontario. Policies and publications [Internet]. Toronto (ON): CPSO; 2018. Available: www.cpso.on.ca/Policies-Publications (accessed 2018 Aug. 13).

AUTHOR RESPONSE

We thank Drs. Byrick and Pitt for their thoughtful letter regarding our recent publication on the topic of propofol administration by nonanesthesiologists in endoscopy. We hope to address a number of excellent discussion points they have brought forward.

In our systematic review and meta-analysis, we searched for studies that compared patient outcomes during endoscopic procedures when propofol was administered by anesthesiologists versus endoscopists. With the intent of capturing a broad population, our inclusion criteria were not restricted to procedures performed inside or outside of hospitals. Despite our initial intent, only one of five studies included patients who received endoscopies outside of a hospital facility,¹ therefore our conclusions are not easily generalizable across settings. It should be noted that we identified very low airway intervention rates, which speaks to the fact that propofol is generally a safe sedative agent, particularly in patients with low comorbidities (i.e., American Society of Anesthesiologists score of I or II). The need for an airway intervention during sedation should not be primarily affected by the facility in which the endoscopy took place, but rather by the patient's comorbidities, the nature of the procedure and the person administering the sedation. For this reason, it is essential that patients who may require advanced anesthetic expertise are identified with the use of validated screening tools. We agree that it is of paramount importance that all facilities, whether inside or outside of hospitals, adhere to local standards of patient safety.

One of the challenges of systematic reviews is the lack of uniform reporting across studies. In the studies included in our review, standards of care were not clearly outlined. Having included studies from various countries, we expect that there were substantial differences in such standards. As described in our table of included studies, one key standard was whether or not the intervention group had a dedicated health care professional in charge of administering sedation and monitoring the patient's cardiorespiratory status throughout the procedure. This was the case in all but one study.¹ Having such a dedicated individual would align with the recommendations established by the College of Physicians and Surgeons of Ontario.²

A new sedation model in which endoscopists provide sedation during endoscopy would require an extensive implementation strategy. First, input from relevant stakeholders needs to be taken into consideration. For instance, a barrier might be the lack of endoscopist buy-in without appropriate incentives. Endoscopists would be required to undertake additional training on managing cardiorespiratory complications. There would also be an increased demand for allied health professionals (i.e., anesthesia assistants, registered nurses) with advanced life support training. A curriculum for

health care professionals to be involved with sedation has already been established by the European Society for Gastrointestinal Endoscopy, which could provide a foundation on the implementation of endoscopist-administered propofol.³ Countries like Canada could also learn from and improve upon the experiences of some European countries where the task of propofol sedation has largely shifted to the endoscopist.^{4,5}

As with any new health system arrangement, there must be an ongoing audit of patient outcomes and worker satisfaction. The costs of such implementation (i.e., education and training, reallocation of health care professionals) need to be weighed against the benefits (i.e., improved access to propofol sedation). Our systematic review serves to highlight that there is low-quality evidence that endoscopists may safely administer propofol to a carefully selected patient population.

Julian F. Daza, MD; Carolyn M. Tan, MD; Ilun Yang, MD

Affiliations: From the Department of Surgery, University of Toronto, Toronto, Ont. (Daza); the Department of Medicine, University of Toronto, Toronto, Ont. (Tan); and the Department of Surgery, McMaster University, Hamilton, Ont. (Yang).

Competing interests: None declared.

DOI: 10.1503/cjs.1861502

References

1. Vargo JJ, Holub JL, Faigel DO et al. Risk factors for cardiopulmonary events during propofol-mediated upper endoscopy and colonoscopy. *Aliment Pharmacol Ther* 2006; 24:955-63.
2. College of Physicians and Surgeons of Ontario. Out-of-Hospital Premises Inspection Program. Toronto (ON): CPSO; 2018. Available: www.cpso.on.ca/CPSO-Members/Out-of-Hospital-Premises-Inspection-Program (accessed 2018 Aug. 30).
3. Dumonceau JM, Riphaut A, Beilenhoff U, et al. European curriculum for sedation training in gastrointestinal endoscopy: position statement of the European Society of Gastrointestinal Endoscopy (ESGE) and European Society of Gastroenterology and Endoscopy Nurses and Associates (ESGENA). *Endoscopy* 2013;45:496-504.
4. Riphaut A, Geist F, Wehrmann T et al. Endoscopic sedation and monitoring practice in Germany: re-evaluation from the first nationwide survey 3 years after the implementation of an evidence and consent based national guideline. *Z Gastroenterol* 2013;51:1082-8.
5. Heuss LT, Froehlich F, Beglinger C et al. Nonanesthesiologist-administered propofol sedation: from the exception to standard practice. Sedation and monitoring trends over 20 years. *Endoscopy* 2012;44:504-11.