

Appendix 1 to Hawel J, McFadgen H, Stewart R, et al. Interventional radiology-assisted transgastric endoscopic drainage of peripancreatic fluid collections. *Can J Surg* 2020.

DOI: 10.1503/cjs.003019

© Joule Inc.

Online appendices are unedited and posted as supplied by the authors.

Appendix 1. PFC Characteristics.

	Mean (SD)	Median (min, max)
Size (cm)	12.70 (3.31)	13.0 (5, 18.3)
	Frequency (%)	
Classification		
Pseudocyst	20 (51.3)	
WON	19 (48.7)	
Infected	20 (51.3)	
Not infected	19 (48.7)	
Etiology		
Acute pancreatitis	32 (82.0)	
Chronic pancreatitis	1 (2.6)	
Acute-on-chronic pancreatitis	1 (2.6)	
Post-op	4 (10.2)	
Post-op pancreatitis	1 (2.6)	
Location		
Head	3 (7.6)	
Body	14 (35.9)	
Tail	4 (10.3)	
Combination	14 (35.9)	
Head & body	2 (14.3)	
Body & tail	10 (71.4)	
Head & Tail	1 (7.1)	
Head, Body, Tail	1 (7.1)	
Not specified	4 (10.3)	
Symptoms		
Pain	30 (76.9)	
Isolated pain	20 (66.7)	
Pain and GOO	3 (10.0)	
Pain and sepsis	1 (3.3)	
Pain and “other”	6 (20.0)	
GOO	8 (20.5)	
Asymptomatic	1 (2.6)	

Appendix 1 to Hawel J, McFadgen H, Stewart R, et al. Interventional radiology-assisted transgastric endoscopic drainage of peripancreatic fluid collections. *Can J Surg* 2020.

DOI: 10.1503/cjs.003019

© Joule Inc.

Online appendices are unedited and posted as supplied by the authors.

Appendix 2. Summary of Complications (n=7)

Classification	Infected vs Sterile	Clavien-Dindo Classification	Details
Pseudocyst	Sterile	Grade IIIb	POD 1: Free air in the abdomen requiring exploratory laparoscopy, no findings at laparoscopy.
Pseudocyst	Infected	Grade IIIb	POD 14: Readmitted for EGD to remove and replace migrated stent. PFC persistent, now infected.
Pseudocyst	Sterile	Grade IIIb	POD 8: Admitted to ICU with sepsis, bacteremia secondary to dislodged stent and persistent PFC. Repeat endoscopy for removal and replacement of stent.
WON	Sterile	Grade IVb	POD 12: Admitted to ICU for resuscitation from massive UGIB from gastrotomy and eroded splenic artery.
Pseudocyst	Sterile	Grade IIIb	POD 11: admitted to ICU w/ sepsis and AKI secondary to blocked stent and persistent, now infected pseudocyst.
Pseudocyst	Sterile	Grade IIIa	Intra-op from IR portion: pneumothorax, chest tube placed in right pleural space
Pseudocyst	Infected	Grade V	POD 49: Admit w/ GIB, no clear source. POD 56: Transferred ICU w/ sepsis secondary to massive retroperitoneal abscess. POD57: died.

POD: postoperative day, EGD: esophagogastroduodenoscopy, ICU: intensive care unit, UGIB: upper gastrointestinal bleed, AKI: acute kidney injury, GIB: gastrointestinal bleed