

Severe street and mountain bicycling injuries in adults: a comparison of the incidence, risk factors and injury patterns over 14 years

Derek J. Roberts, MD*†
 Jean-Francois Ouellet, MD*§
 Francis R. Sutherland, MD*
 Andrew W. Kirkpatrick, MD,
 MHSc*†§
 Rohan N. Lall, MD*§
 Chad G. Ball, MD, MSc*§

From the Departments of *Surgery, †Community Health Sciences, and ‡Critical Care Medicine as well as the §Regional Trauma Program, University of Calgary and the Foothills Medical Centre, Calgary, Alta.

This study was presented at the 2011 Canadian Surgery Forum in London, Ont. on Sept. 16, 2011.

Accepted for publication
 May 22, 2012

Correspondence to:

C.G. Ball
 Trauma and Acute Care Surgery
 Hepatobiliary and Pancreatic Surgery
 Departments of Surgery and Oncology
 University of Calgary and Foothills
 Medical Centre
 1403–29th Street NW
 Calgary AB T2N 2T9
 ball.chad@gmail.com

DOI: 10.1503/cjs.027411

Background: Street and mountain bicycling are popular recreational activities and prevalent modes of transportation with the potential for severe injury. The purpose of this investigation was to compare the incidence, risk factors and injury patterns among adults with severe street versus mountain bicycling injuries.

Methods: We conducted a retrospective cohort study using the Southern Alberta Trauma Database of all adults who were severely injured (injury severity score [ISS] ≥ 12) while street or mountain bicycling between Apr. 1, 1995, and Mar. 31, 2009.

Results: Among 11 772 severely injured patients, 258 (2.2%) were injured (mean ISS 17, hospital stay 6 d, mortality 7%) while street ($n = 209$) or mountain bicycling ($n = 49$). Street cyclists were often injured after being struck by a motor vehicle, whereas mountain bikers were frequently injured after faulty jump attempts, bike tricks and falls (cliffs, roadsides, embankments). Mountain cyclists were admitted more often on weekends than weekdays (61.2% v. 45.0%, $p = 0.040$). Injury patterns were similar for both cohorts (all $p > 0.05$), with trauma to the head (67.4%), extremities (38.4%), chest (34.1%), face (26.0%) and abdomen (10.1%) being common. Spinal injuries, however, were more frequent among mountain cyclists (65.3% v. 41.1%, $p = 0.003$). Surgical intervention was required in 33.3% of patients (9.7% open reduction internal fixation, 7.8% spinal fixation, 7.0% craniotomy, 5.8% facial repair and 2.7% laparotomy).

Conclusion: With the exception of spine injuries, severely injured cyclists display similar patterns of injury and comparable outcomes, regardless of style (street v. mountain). Helmets and thoracic protection should be advocated for injury prevention.

Contexte : Le vélo de ville et le vélo de montagne sont des activités récréatives et des modes de transport populaires très utilisés, qui comportent un risque de blessures graves. Le but de la présente étude était de comparer l'incidence des blessures, les facteurs de risque et les types de blessures les plus fréquents chez les adultes victimes d'accidents impliquant l'utilisation de la bicyclette en ville et hors-piste.

Méthodes : Nous avons procédé à une étude de cohorte rétrospective à partir de la base de données de traumatologie du Sud de l'Alberta, en regroupant tous les adultes qui ont été victimes d'une blessure grave (indice de gravité de la blessure [IGB] ≥ 12), alors qu'ils circulaient à vélo en ville ou hors-piste entre le 1^{er} avril 1995 et le 31 mars 2009.

Résultats : Parmi les 11 772 patients blessés gravement, 258 (2,2 %) l'ont été (IGB moyen 17, séjour hospitalier 6 jours, mortalité 7 %) alors qu'ils circulaient à bicyclette en ville ($n = 209$) ou hors-piste ($n = 49$). Les cyclistes qui roulent en ville sont souvent victimes de collision avec des automobiles, tandis que les adeptes du vélo de montagne se blessent souvent lors de tentatives de sauts ou d'acrobaties infructueuses et de chutes (escarpements, accotements, talus). Les adeptes du vélo de montagne ont été plus souvent admis les fins de semaine que les jours de semaine (61,2 % c. 45,0 %, $p = 0,040$). Les types de blessures étaient similaires dans les 2 groupes (tous $p > 0,05$), les traumatismes crâniens (67,4 %), les blessures aux extrémités (38,4 %), à la poitrine (34,1 %), au visage (26,0 %) et à l'abdomen (10,1 %) ayant été les plus fréquents. Les lésions médullaires ont toutefois été plus fréquentes chez les adeptes du vélo de montagne (65,3 % c. 41,1 %, $p = 0,003$). Une intervention chirurgicale a été nécessaire chez 33,3 % des patients (9,7 % pour fixation interne par réduction chirurgicale, 7,8 % pour une fixation du rachis, 7,0 % pour une craniotomie, 5,8 % pour réparation faciale et 2,7 % pour laparotomie).

Conclusion : À l'exception des blessures à la colonne vertébrale, les cyclistes gravement blessés présentent des types de blessures similaires et des résultats comparables, indépendamment du type de vélo qu'ils pratiquent (ville c. montagne). Il faut promouvoir le port de casques et de plastrons de protection pour prévenir les blessures.

Street and mountain bicycling are increasingly popular in North America, with Canadian biking rates exceeding those of the United States.¹ In southern Alberta, both bicycling styles are prevalent, given the region's climate and biking infrastructure.^{1,2} Chinook winds and reduced snowfall result in improved roadway conditions and warmer temperatures year-round.² Moreover, about 370 km of bicycling lanes and pathways are available within the city of Calgary, resulting in the largest network of bike paths per 100 000 population in Canada.^{1,2} Surrounding regions, such as Kananaskis and the Rocky Mountains, further contribute to making the Calgary area one of the busiest biking environments in Canada.

Unfortunately, street and mountain bicycling injuries can be severe and even fatal.³⁻⁷ Absolute injury event rates per kilometer for street cycling range from 26 to 68 times higher than those for motor vehicle travel.⁸ In British Columbia, major trauma related to street cycling occurs at an annual incidence of 49.8 per 100 000 (2.9% of all trauma admissions), with most deaths occurring as a result of neurotrauma.⁴ Conversely, mountain biking injuries accounted for only 1.6% of all severe trauma patient admissions in this area from 1992 to 2002 (about 1-6 injuries per 100 000 patients per year).⁵

Although observational studies of bicycling injuries are common,^{3-7,9-17} to our knowledge, our study is the first to compare the overall major trauma epidemiology of street and mountain bicycling in detail. Few reports have discussed experience with severe bicycling-related multisystem injuries that require emergent surgical intervention or intensive care unit (ICU) admission. Our objective was to compare the incidence, risk factors and injury patterns of street versus mountain bicycling-related major trauma over a 14-year period.

METHODS

Using the Southern Alberta Trauma Database, we performed a descriptive retrospective cohort study comparing all adults (≥ 16 yr) who were severely injured (injury severity score [ISS] ≥ 12) while street or mountain bicycling between Apr. 1, 1995, and Mar. 31, 2009. The registry contains prospectively collected data on all trauma patients admitted to the Foothills Medical Centre (FMC), regardless of their length of stay. The FMC is an adult, level 1 tertiary care trauma hospital responsible for all major injuries in southern Alberta, southwestern Saskatchewan and southeastern British Columbia. The University of Calgary Conjoint Health Research Ethics Board approved the study.

The following data were extracted for all severely injured cyclists: demographic characteristics, mechanism of injury, pattern of injury, specific injuries, ISS, mode of transport to hospital, mean patient transport time (i.e., the time from injury to arrival at our trauma centre) and surgical interventions. Although bicycle helmet legislation exists in Alberta and 48.3% of Albertans aged 12 years and

older wear a helmet while cycling,¹⁸ helmet use is poorly documented in the trauma database as it is often not described in patient charts, emergency medical services (EMS) documents and STARS air ambulance reports. Outcomes included discharge status (alive or dead) and length of hospital and ICU stay. To determine whether there were temporal or seasonal influences on the rate of bicycling-related injury, we also examined the incident date, season and time of day (early morning, 12:01 am-8:00 am; morning/afternoon, 8:01 am-4:00 pm; evening, 4:01 pm-8:00 pm; or night 8:01 pm-12:00 am), as well as whether the injury occurred on a weekend (between 5:00 pm on Friday and 8:00 am on Monday) or on a weekday.

Statistical analysis

We conducted comparative analyses using Stata software, version 8.0 (Stata Corp.). Normally distributed variables are reported as means and standard deviation (SD), and non-normally distributed data are reported as medians with interquartile ranges (IQRs). We compared means using Student *t* tests and medians using the Mann-Whitney *U* test. Differences in proportions for categorical data required Fisher exact or χ^2 tests. We considered results to be significant at $p < 0.05$, 2-tailed.

RESULTS

Over the 14-year study period, 11 772 patients required admission to the FMC for management of injuries with an ISS of 12 or greater. Of these, 258 (2.2%) patients were injured (median ISS 17 [IQR 16-25]) during street cycling ($n = 209$) or mountain biking ($n = 49$; Table 1). The clinical characteristics of the 2 cohorts were similar, with the exception that mountain bikers were significantly younger than street cyclists (28 v. 43 yr, $p < 0.001$). Most patients were male (86.4%), and all were operators of the bicycle. Discharge destinations included home ($n = 189$), rehabilitation facilities ($n = 36$) and acute care facilities ($n = 11$). Five patients left against medical advice, and the remaining patients died ($n = 17$).

Street cycling was associated with a significantly higher overall rate of admission for severe injury than mountain bicycling (1.8% v. 0.42%, $p < 0.001$). Although fluctuations occurred across calendar years, there was no significant difference in the yearly rate of admission for either style of bicycling (Fig. 1). The median overall hospital stay for admitted patients was 6 (IQR 3-12) days and was similar between the 2 cohorts. Admission to the ICU was required for 53 (25.4%) street and 15 (30.6%) mountain cyclists (both $p > 0.05$; similar median ICU stay of 3 [IQR 1-3] d).

The time of day and season during which most bicycling-related trauma occurred was similar for street and mountain cyclists, with most injuries occurring in the morning or afternoon (24.0%) and in summer (53.9%) or spring

(29.5%). A trend toward a greater number of bicycling injuries in the summer (53.9%) versus other seasons was observed ($p = 0.09$; Table 2). When compared with street cyclists, mountain bikers were admitted to hospital more commonly on weekends (61.2% v. 45.0%, $p = 0.040$).

Most injured cyclists were transported to hospital via ground (74.4%) versus air ambulance (16.7%) or private vehicle (7.9%; Table 1). More mountain than street cyclists were transported by air ambulance (36.7% v. 11.9%, $p < 0.001$). Conversely, transport by ground was significantly more common for street than mountain cyclists (79.0% v. 55.1%, $p < 0.001$). Accordingly, the mean transport time was significantly longer for mountain (43 [IQR 22–89] min) than street cyclists (19 [IQR 12–60] min; $p = 0.020$).

Cyclists' mechanisms of injury are presented in Table 3. Major injury among street cyclists occurred commonly after being hit by a motor vehicle (27.8%), whereas mountain bikers were often severely injured after faulty jump attempts or bike tricks (20.4%) and falls off cliffs, roadsides and embankments (16.3%; all $p < 0.05$).

These injury mechanisms resulted in largely similar anatomic injury patterns across both groups (all $p > 0.05$), with trauma to the head (67.4%), extremities (38.4%), chest (34.1%), face (26.0%) and abdomen (10.1%) being common. Spinal injuries, however, were significantly more frequent among mountain cyclists (65.3% v. 41.1%, $p = 0.003$; Fig. 2). Specific injuries are described in detail in Table 4.

Surgical intervention was required in 33.3% of patients and was similar between groups (Table 5). Extremity open reduction internal fixation was most common (9.7%), followed by spinal fixation (7.8%), craniotomy (7.0%), facial fracture repair (5.8%) and laparotomy (2.7%). Indications for laparotomy included renal laceration ($n = 1$: American Association for the Surgery of Trauma [AAST] grade III), splenic injuries ($n = 4$: 3 AAST grade V, 1 grade III), duodenal laceration ($n = 1$: AAST grade III) and pelvic fracture ($n = 1$) requiring extraperitoneal packing after pelvic angiobolization.

A total of 240 (93.0%) patients were discharged alive. Of the 17 (6.6%) cyclists (16 street cyclists and 1 mountain biker) who died from their injuries, 14 had a severe traumatic brain injury (3 also had an extensive extra- or intracranial cerebrovascular artery dissection and another had an associated complete atlantoaxial dislocation), 1 sustained a complete spinal cord injury at the C1/C2 level and 2 succumbed to a combined blunt thoracic aortic injury, open book pelvic fracture and internal iliac artery laceration or an AAST grade II liver and grade III renal injury.

DISCUSSION

Although several investigations have delineated the injury




Fig. 1. Annual number of trauma centre admissions for severe street and mountain bicycling injuries between 1996 and 2008. The years 1995 and 2009 were excluded, as those calendar years were incomplete in this investigation. No significant difference in annual admission rate was observed between street and mountain bicycling or between any 2 years for either style of bicycling.

Table 1. Characteristics of the severely injured cyclists included in our study

Characteristic	Overall, $n = 258$	Street cyclists, $n = 209$	Mountain bikers, $n = 49$	p value
Age, median (IQR) yr	39 (27–53)	43 (0–55)	28 (21–35)	< 0.001
Male sex, no. (%)	223 (86.4)	180 (86.1)	43 (87.8)	0.09
Mode of transportation to hospital, no. (%)				
Private vehicle	20 (7.9)	16 (7.7)	4 (8.2)	> 0.99
Ground ambulance	192 (74.4)	165 (79.0)	27 (55.1)	< 0.001
Air ambulance	43 (16.7)	25 (11.9)	18 (36.7)	< 0.001
Transportation time, median (IQR) min	23 (13–66)	19 (12–60)	43 (22–89)	0.02
ICU admission required, no. (%)	68 (26.4)	53 (25.4)	15 (30.6)	0.47
Length of ICU stay, median (IQR) d	3 (1–3)	3 (1–12)	4 (3–10)	0.79
Length of hospital stay, median (IQR) d	6 (3–12)	6 (3–13)	5 (4–9)	0.79
Death, no. (%)	17 (6.6)	16 (7.7)	1 (2.0)	0.21
ISS, median (IQR)	17 (16–25)	17 (16–25)	17 (15–25)	0.70
GCS score, median (IQR)	15 (14–15)	15 (14–15)	15 (15–15)	> 0.99

GCS = Glasgow Coma Scale; ICU = intensive care unit; IQR = interquartile range; ISS = injury severity score; SD = standard deviation.

epidemiology of less severely injured cyclists,^{9,10,12-17,19} this retrospective cohort study analyzed only patients with severe multisystem injuries (ISS \geq 12). This was affirmed by an overall median ISS of 17, ICU admission rate of 26.4%, surgical intervention rate of 33.3% and mortality of 6.6%. The associated anatomic injury patterns and trauma epidemiology of these patients appear to differ markedly from those of less severely injured cyclists.

In addition to variances in injury mechanisms between street and mountain cyclists, we identified several risk factors for severe cycling-related trauma. Overall, injury was more frequent among street cyclists, possibly because street cycling is an extremely prevalent, year-round mode of transportation in southern Alberta. This finding could also have resulted from some mountain bikers receiving care at hospitals in surrounding mountain regions rather than at the FMC. However, we believe this to be an unlikely ex-

planation, as the FMC is the receiving hospital for major trauma for southern Alberta, southwestern Saskatchewan and southeastern British Columbia. Although bicycle-level falls were the most common injury mechanism, street cyclists were commonly injured after being hit by a motor vehicle, whereas mountain bikers were often injured after faulty jump attempts, bike tricks and falls off cliffs, roadsides and embankments. As expected, most injuries occurred in the morning or afternoon and in summer or spring. Compared with street cyclists, mountain bikers were also more commonly admitted on weekends, likely because a commute to the mountains is often desired.

As reported in other studies,³⁻⁷ traumatic brain injury was common, occurring in 67% of severely injured street and mountain cyclists. Up to 7% of these patients required an emergent craniotomy. Interestingly, this injury rate is identical to that reported by Guichon and Myles¹¹ in their

Table 2. Individual or environmental characteristics associated with severe bicycling trauma

Characteristic	Group; no. (%)			p value
	Overall, n = 258	Street cyclists, n = 209	Mountain bikers, n = 49	
Time of year				0.09
Winter (Dec. 21 to Mar. 19)	11 (4.3)	10 (4.8)	1 (2.0)	
Spring (Mar. 20 to June 20)	76 (29.5)	62 (29.7)	14 (28.6)	
Summer (June 21 to Sept. 22)	139 (53.9)	106 (50.7)	33 (67.3)	
Fall (Sept. 23 to Dec. 20)	28 (10.9)	27 (12.9)	1 (2.0)	
Unknown	4 (1.6)	4 (1.9)	0 (0)	
Time of the week				0.029
Weekday	134 (51.9)	115 (55.0)	19 (38.8)	
Weekend*	124 (48.1)	94 (45.0)	30 (61.2)	
Time of the day				0.17
Early morning (12:01 am–8:00 am)	18 (7.0)	16 (7.7)	2 (4.1)	
Morning/afternoon (8:01 am–4:00 pm)	62 (24.0)	46 (22.0)	16 (32.7)	
Evening (4:01 pm–8:00 pm)	39 (15.1)	36 (17.2)	3 (6.1)	
Night (8:01 pm–12:00 am)	16 (6.2)	12 (5.7)	4 (8.2)	
Unknown	123 (47.7)	99 (47.4)	24 (49.0)	
Ethanol usage	5 (1.9)	5 (2.4)	0 (0)	0.59

*A weekend was defined as the time period between 5:00 pm on Friday and 8:00 am on Monday.

Table 3. Specific mechanism of injury distribution*

Mechanism	Group; no. (%)			p value
	Overall, n = 258	Street cyclists, n = 209	Mountain bikers, n = 49	
Fell off bicycle	120 (46.5)	93 (44.5)	27 (55.1)	0.20
Attempted to avoid a person, animal or object	5 (1.9)	5 (2.4)	0 (0)	
Rode down a hill at high speed	19 (7.4)	19 (9.1)	0 (0)	
Hit a speed bump	2 (0.7)	2 (0.9)	0 (0)	
Lost balance, speed-related or other cause	94 (36.4)	67 (32.0)	27 (55.1)	
Fell while attempting a jump or trick	29 (11.2)	19 (9.1)	10 (20.4)	0.041
Veered and fell off cliff, roadside or embankment	14 (5.4)	6 (2.9)	8 (16.3)	0.001
Collided with a person, animal or object other than a motor vehicle	28 (10.9)	25 (12.0)	3 (6.1)	0.31
Collided with a parked automobile	7 (2.7)	7 (3.3)	0 (0)	
Hit by a motor vehicle while bicycling	58 (22.5)	58 (27.8)	0 (0)	
Hit by a commuter train while crossing tracks	1 (0.39)	1 (0.48)	0 (0)	

*p values not afforded when event rates too small to compare or when comparisons would not be meaningful (e.g., collision with parked car while mountain biking).

pioneering 1975 survey of bicycling-related trauma in Calgary. In accordance with a National Trauma Databank study, we identified cerebral contusion (22%), subarachnoid hemorrhage (22%) and subdural hematoma (26%) as the most frequent types of brain injury.⁶ Epidural hematomas




Fig. 2. Anatomic injury distribution for the 258 severely injured street and mountain cyclists. Injury rates were all similar between both groups (all $p > 0.05$).

were also common among severely injured cyclists (9%). This may be owing to the predominance of young men in our study, who are known to possess weak attachments between the dura and inner table of the skull.⁶ They are also more likely to participate in an aggressive, high-speed riding technique.⁶ Although helmet use reduces the odds of head injury by 63%–88%,²⁰ the frequency and effectiveness of this intervention could not be determined in our cohort, as helmet usage was poorly documented.

Spinal injury was also common (45.7%) and resulted in concomitant injury to the spinal cord in 8.5% of patients. The rate of spinal trauma was significantly higher among mountain than street cyclists (65.3% v. 41.1%, $p = 0.003$), with fractures to the cervical and thoracolumbar spine being equally frequent.⁵ A potential mechanism for concomitant head and cervical spine injury in mountain bikers has previously been postulated to involve a fall over the handlebars, resulting in a blow to the top of the head and cervical flexion or hyperextension.^{5,21} This mechanism may easily have occurred in this study, given that so many mountain bikers were observed to fall off cliffs, roadsides and embankments.

In addition to head and spinal trauma, many street and mountain cyclists also presented with extremity (38%), thoracic (34%) and abdominal injuries (10%). Although the orthopedic injury rate approximated those observed in previous investigations,^{4-6,22,23} the frequency of thoracic

Table 4. Specific bicycling-related injuries*

Specific injury	Group; no. (%)			p value
	Overall, n = 258	Street cyclists, n = 209	Mountain bikers, n = 49	
Head	174 (67.4)	146 (69.9)	28 (57.1)	0.09
Cerebral contusion	56 (21.7)	46 (22.0)	10 (20.4)	
Concussion	43 (16.7)	37 (17.7)	6 (12.2)	
Diffuse axonal injury	16 (6.2)	11 (5.3)	5 (10.2)	
Epidural hematoma	22 (8.5)	20 (9.6)	2 (4.1)	
Subarachnoid hemorrhage	56 (21.7)	46 (22.0)	10 (20.4)	
Subdural hematoma	67 (26.0)	58 (27.8)	9 (18.4)	
Spine	118 (45.7)	86 (41.1)	32 (65.3)	0.003
Cervical fracture	50 (19.4)	37 (17.7)	13 (26.5)	
Thoracolumbar fracture	46 (17.8)	33 (15.8)	13 (26.5)	
Cord injury	22 (8.5)	16 (7.7)	6 (12.2)	
Thorax	88 (34.1)	73 (34.9)	15 (30.6)	0.62
Hemo- and/or pneumothorax	51 (19.8)	38 (18.2)	13 (26.5)	
Pulmonary contusion(s)	29 (11.2)	20 (9.6)	9 (18.4)	
Rib fracture(s)	60 (23.3)	52 (24.9)	8 (16.3)	
Abdomen	26 (10.1)	22 (10.5)	4 (8.2)	0.79
Kidney	7 (2.7)	6 (2.9)	1 (2.0)	
Liver	5 (1.9)	4 (1.9)	1 (2.0)	
Small bowel	2 (0.78)	1 (0.48)	1 (2.0)	
Spleen	14 (5.4)	12 (5.7)	2 (4.1)	
Bony pelvis	17 (6.6)	15 (7.2)	2 (4.1)	0.75
Extremity	99 (38.4)	77 (36.8)	20 (40.8)	0.63
Upper fractures	33 (12.8)	25 (12.0)	8 (16.3)	0.48
Lower fractures	32 (12.4)	21 (10.0)	0 (0)	0.001

*The sum of patients in each category does not always equal the total as many patients had multiple injuries.

injuries was significantly higher. This most commonly comprised rib fractures, pulmonary contusions and hemothoraces. Similar to the rate of chest trauma among horseback riders in our area,²⁴ chest trauma may be underappreciated in severely injured cyclists, highlighting the potential importance of thoracic protection. While reports of subcapsular and central liver hematomas were common before the elimination of upright bicycle handlebars,^{14,25} the liver was uncommonly injured (1.9%) compared with the spleen (5.4%) among our cohorts.

A total of 17 patients died over the 14-year study period. These deaths were primarily a result of traumatic brain injury, cervical spinal cord injury and/or physiologic exhaustion from ongoing hemorrhage. This is supported by Fife and colleagues,²² who observed that the causes of bicycling-related deaths included early (hemorrhage, cerebral edema and aspiration) and late (pneumonia and pulmonary embolism) etiologies. Age, ISS and alcohol consumption have also been found previously to be associated with increased mortality among injured cyclists.^{3,6,22}

Limitations

Although, to our knowledge, this study is the only comparison of major injuries between street and mountain cyclists, it has several limitations. Given the retrospective design, the possibility of bias cannot be eliminated. Moreover, the most glaring omission was poor documentation of helmet usage, which was unavailable for 85% of all patients identified in the trauma database. Although 48.3% of Albertans older than 12 years wear a helmet while cycling,¹⁸ helmet use is frequently not described in

patient charts, EMS documents and STARS air ambulance reports. All other variables were well described. Finally, we were unable to characterize individual cyclist skill level, which may impact both injury rates and severity.

CONCLUSION

The overall rate of trauma centre admission for severe bicycling-related injuries in southern Alberta over 14 years was 2.2%. This was composed primarily of street cyclists. Despite differing style (street v. mountain), severely injured cyclists display largely similar injury patterns with comparable outcomes. While more mountain bikers are injured on weekends, they require surgical interventions at the same rate as street cyclists. Helmet and thoracic protection should be advocated as injury prevention measures.

Acknowledgment: We thank Mrs. Corina Tiruta for assistance with statistical analyses.

Competing interests: None declared.

Contributors: D.J. Roberts, J.-F. Ouellet, F.R. Sutherland and C.G. Ball designed the study. D.J. Roberts, F.R. Sutherland, R.N. Lall and C.G. Ball acquired the data. D.J. Roberts, J.-F. Ouellet, F.R. Sutherland, A.W. Kirkpatrick and C.G. Ball analyzed the data. D.J. Roberts and C.G. Ball wrote the article. All authors reviewed the article and approved its publication.

References

1. Pucher J, Buehler R. Why Canadians cycle more than Americans: a comparative analysis of bicycling trends and policies. *Transp Policy* 2006;13:265-79.
2. Calgary downtown commuter cyclist survey report. Calgary (AB): The City of Calgary; 2006. Available: www.calgary.ca/Transportation

Table 5. Surgical interventions performed and associated indications*

Intervention	Group; no. (%)			p value
	Overall, n = 258	Cyclists, n = 209	Mountain bikers, n = 49	
Craniotomy or craniectomy for evacuation of hematoma	18 (7.0)	17 (8.1)	1 (2.0)	0.13
Facial procedure for fracture(s)	15 (5.8)	11 (5.3)	4 (8.2)	0.43
Spinal procedure for fracture(s) or instability	20 (7.8)	14 (6.7)	6 (12.2)	0.19
Thoracotomy for	2 (0.8)	1 (0.5)	1 (2.0)	
Hemorrhaging LLL laceration and pulmonary artery pseudoaneurysm	1 (0.4)	0 (0)	1 (2.0)	
Intercostal artery injury				
Laparotomy for:	7 (2.7)	7 (3.3)	0 (0)	0.19
AAST grade II liver and grade III renal injury	1 (0.4)	1 (0.5)	0 (0)	
Splenic injury	1 (0.4)	1 (0.5)	0 (0)	
AAST grade III	3 (1.2)	3 (1.4)	0 (0)	
AAST grade V	1 (0.4)	1 (0.5)	0 (0)	
Duodenal perforation	1 (0.4)	1 (0.5)	0 (0)	
Extraperitoneal packing of bleeding pelvic fracture				
Extremity open reduction internal fixation	25 (9.7)	23 (11.0)	2 (4.1)	0.14
Vascular repair for brachial artery transection	1 (0.4)	0 (0)	1 (2.0)	
Repair of blunt tracheolaryngeal injury	1 (0.4)	1 (0.5)	0 (0)	
Wound suturing and/or debridement alone	5 (1.9)	5 (2.4)	0 (0)	
None	172 (66.7)	138 (66.0)	34 (69.4)	0.65

AAST = American Association for the Surgery of Trauma; LLL = left lower lobe of the lung.
*p values not afforded when event rates too small to compare.

- /TP/Documents/transportation_solutions/downtown_cyclist_survey_2007_p1of4.pdf (accessed 2011 Jul. 1).
3. Frank E, Frankel P, Mullins RJ, et al. Injuries resulting from bicycle collisions. *Acad Emerg Med* 1995;2:200-3.
 4. Konkin DE, Garraway N, Hameed SM, et al. Population-based analysis of severe injuries from nonmotorized wheeled vehicles. *Am J Surg* 2006;191:615-8.
 5. Kim PT, Jangra D, Ritchie AH, et al. Mountain biking injuries requiring trauma center admission: a 10-year regional trauma system experience. *J Trauma* 2006;60:312-5.
 6. Lustenberger T, Inaba K, Talving P, et al. Bicyclists injured by automobiles: relationship of age to injury type and severity — a National Trauma Databank analysis. *J Trauma* 2010;69:1120-5.
 7. Sacks JJ, Holmgren P, Smith SM, et al. Bicycle-associated head injuries and deaths in the United States from 1984 through 1988. How many are preventable? *JAMA* 1991;266:3016-8.
 8. Aultman-Hall L, Kaltenecker MG. Toronto bicycle commuter safety rates. *Accid Anal Prev* 1999;31:675-86.
 9. Gassner RJ, Hackl W, Tuli T, et al. Differential profile of facial injuries among mountainbikers compared with bicyclists. *J Trauma* 1999;47:50-4.
 10. Gaulrapp H, Weber A, Rosemeyer B. Injuries in mountain biking. *Knee Surg Sports Traumatol Arthrosc* 2001;9:48-53.
 11. Guichon DM, Myles ST. Bicycle injuries: one-year sample in Calgary. *J Trauma* 1975;15:504-6.
 12. Jeys LM, Cribb G, Toms AD, et al. Mountain biking injuries in rural England. *Br J Sports Med* 2001;35:197-9.
 13. Maimaris C, Summers CL, Browning C, et al. Injury patterns in cyclists attending an accident and emergency department: a comparison of helmet wearers and non-wearers. *BMJ* 1994;308:1537-40.
 14. Nehoda H, Hochleitner BW, Hourmont K, et al. Central liver hematomas caused by mountain-bike crashes. *Injury* 2001;32:285-7.
 15. Nicaj L, Stayton C, Mandel-Ricci J, et al. Bicyclist fatalities in New York City: 1996–2005. *Traffic Inj Prev* 2009;10:157-61.
 16. Rivara FP, Thompson DC, Thompson RS, et al. Injuries involving off-road cycling. *J Fam Pract* 1997;44:481-5.
 17. Rowe BH, Rowe AM, Bota GW. Bicyclist and environmental factors associated with fatal bicycle-related trauma in Ontario. *CMAJ* 1995; 152:45-53.
 18. Bicycle helmet use, 2009. In: *Canadian Statistics* [website of Statistics Canada]. Available: www.statcan.gc.ca/pub/82-625-x/2010002/article/11274-eng.htm (accessed 2012 Apr. 3).
 19. Spaite DW, Murphy M, Criss EA, et al. A prospective analysis of injury severity among helmeted and nonhelmeted bicyclists involved in collisions with motor vehicles. *J Trauma* 1991;31:1510-6.
 20. Thompson DC, Rivara FP, Thompson R. Helmets for preventing head and facial injuries in bicyclists. *Cochrane Database Syst Rev* 2000;(2): CD001855.
 21. Carmont MR. Mountain biking injuries: a review. *Br Med Bull* 2008; 85:101-12.
 22. Fife D, Davis J, Tate L, et al. Fatal injuries to bicyclists: the experience of Dade County, Florida. *J Trauma* 1983;23:745-55.
 23. Boström L, Nilsson B. A review of serious injuries and deaths from bicycle accidents in Sweden from 1987 to 1994. *J Trauma* 2001;50: 900-7.
 24. Ball CG, Ball JE, Kirkpatrick AW, et al. Equestrian injuries: incidence, injury patterns, and risk factors for 10 years of major traumatic injuries. *Am J Surg* 2007;193:636-40.
 25. Nehoda H, Hochleitner BW. Subcapsular liver haematomas caused by bar ends in mountain-bike crashes. *Lancet* 1998;351:342.