

Laparoscopic bariatric surgery can be performed safely in secondary health care centres with a dedicated service corridor to an affiliated tertiary health care centre

Nicolas Christou, MD, PhD

From the Department of Surgery, McGill University, and the Section of Bariatric Surgery, Division of General Surgery, McGill University Health Centre, Montréal, Que.

Presented in part at the Canadian Surgery Forum in London, Ont., Sept. 15–18, 2011.

Accepted for publication July 30, 2012

Correspondence to:

N.V. Christou
McGill University Health Centre
Room S6.24
687 Pine Ave. W
Montréal QC H3A 1A1
Nicolas.christou@mcgill.ca

DOI: 10.1503/cjs.005612

Background: Canada needs to increase capacity for bariatric surgery to reduce the wait for this cost-effective, life-saving surgery. The aim of this study was to test whether laparoscopic bariatric surgery, including gastric bypass, can be delivered safely in secondary health care centres (SHCCs).

Methods: In this prospective cohort study, patients received bariatric surgery at an SHCC that had no intensive care unit but had a dedicated operating room and ward teams and a patient-monitoring environment. Patients with life-threatening complications were transferred to an affiliated tertiary health care centre (THCC) via a dedicated “service corridor.”

Results: In all, 830 patients were treated: 676 at the SHCC and 154 at the THCC. Gastric bypass was performed in 85.4%, gastric band in 11.1% and gastric sleeve in 3.5%. The body mass index (BMI) was significantly higher in the THCC than the SHCC group (mean 54.4 [standard deviation (SD) 9.7] v. 47.5 [SD 7.4]). Obesity-associated diseases were similar between the groups. Major complications occurred in 2.6% of SHCC patients and 1.7% of THCC patients. Seven patients (1%) required direct transfer to the THCC, and all were treated successfully. There were 2 deaths (1.3%) in the THCC and none in the SHCC groups (combined mortality 0.2%). Weight loss was equivalent up to the fourth year of the study.

Conclusion: With proper patient selection, a dedicated health care team and a service corridor to an affiliated THCC, laparoscopic bariatric surgery, including gastric bypass can be performed safely in SHCCs. Further study is needed to determine whether the model can be applied across Canada.

Contexte : Le Canada doit accroître sa capacité en chirurgie bariatrique afin de réduire les temps d'attente pour cette intervention rentable qui sauve des vies. Le but de cette étude était de vérifier si la chirurgie bariatrique laparoscopique, y compris le pontage gastrique laparoscopique, peut se pratiquer en toute sécurité dans les centres hospitaliers de soins de deuxième ligne (CHDL).

Méthodes : Dans cette étude de cohorte prospective, les patients ont subi une chirurgie bariatrique dans un CHDL sans unité de soins intensifs mais pourvu d'un bloc opératoire et d'équipes soignantes spécialisées, ainsi que d'un système de surveillance des patients. Les patients qui ont présenté des complications gravissimes ont été transférés dans un centre hospitalier de soins tertiaires (CHST), par le biais d'un « corridor de service » réservé.

Résultats : En tout, 830 patients ont subi le traitement; 676 au CHDL et 154 au CHST. On a procédé par pontage gastrique chez 85,4 % patients, pose d'un anneau gastrique chez 11,1 % et gastrectomie longitudinale en manchon (gastric sleeve) chez 3,5 % des patients. L'indice de masse corporelle (IMC) était significativement plus élevé dans le groupe transféré au CHST que dans le groupe traité au CHDL (moyenne 54,4 [écart-type (ET) 9,7 c. 47,5 [ET 7,4]). Les maladies associées à l'obésité étaient similaires entre les groupes. Des complications majeures sont survenues chez 2,6 % des patients du CHDL et 1,7 % des patients du CHST. Sept patients (1 %) ont dû être transférés directement au CHST et ils ont tous été traités avec succès. On a déploré 2 décès (1,3 %) dans le groupe du CHST; aucun n'est survenu chez les patients du CHDL (mortalité combinée 0,2 %). La perte de poids a été équivalente jusqu'à la quatrième année de l'étude.

Conclusion : Avec une sélection adéquate des patients, une équipe soignante spécialisée et un corridor de service vers un CHST affilié, il est possible d'effectuer sécuritairement des chirurgies bariatriques laparoscopiques, y compris par pontage gastrique, dans les CHDL. Il faudra approfondir la recherche pour déterminer si ce modèle peut être appliqué partout au Canada.

According to the World Health Organization, obesity is reaching epidemic proportions. Canada is no exception to this epidemic: most Canadians are overweight or obese,¹ and 2% of men and 4% of women are morbidly obese.² Obesity-related death rates are at least on par with deaths related to smoking, and some authors believe that obesity is now the number 1 killer in North America.³

Bariatric surgery is the only treatment that produces substantial, sustained, long-term weight loss in patients with severe obesity.^{4,5} In addition, permanent weight loss through bariatric surgery reduces the relative risk of death by 35%–89%^{6–10} and produces substantial pharmacoeconomic benefits.¹¹ Despite these well-documented findings, bariatric surgery is difficult to access in Canada.^{12,13}

In 2006, a unique pilot project was initiated to determine whether laparoscopic bariatric surgery can be safely performed in smaller hospitals, designated as secondary health care centres (SHCCs), and linked via a dedicated service corridor to a full service tertiary health care centre (THCC). The model was proposed by l'Agence d'évaluation des technologies et des modes d'intervention en santé (AETMIS) in their report to the Quebec Minister of Health and Social Services as a means of increasing bariatric surgery capacity in the province.¹⁴ The present article outlines the findings of this pilot project.

METHODS

This pilot project was initiated following publication of the AETMIS report and a meeting with the Quebec Minister of Health and Social Services at the time. The 534-bed McGill University Health Centre (MUHC), which has more than 40 years of bariatric surgery experience, is fully equipped with an intensive care unit (ICU) and has dialysis capability, was selected as the THCC. The SHCC was the Centre Métropolitain du chirurgie, a fully accredited 17-bed private hospital with a "Specialized Medical Centre" designation from the Ministry of Health and Social Services.¹⁵ The SHCC has no ICU, but has a patient-to-nurse ratio of 1:3 and oxygen saturation monitoring capacity for all patients, essentially providing "step-down unit" care for the patients until discharge. The same surgeon performed all procedures with the same dedicated operating room team, ward nurses and support staff over the duration of the study. Patients with potentially life-threatening complications were transferred to the THCC via a special ambulance using a priori determined protocol (dedicated "service corridor").

All patients met the 1991 National Institutes of Health Consensus Conference guidelines¹⁶ for bariatric surgery: a body mass index (BMI) greater than 35 with associated comorbidities, or a BMI greater than 40. All patients were assessed, including medical, nutritional and psychological assessments, by a multidisciplinary team. Uncontrollable binge-eating disorders required treatment before surgery. All patients were also required to demonstrate an understanding of the surgical procedure they were scheduled to undergo, its mechanism of weight loss, potential long- and short-term complications, dietary and physical activity requirements, and the need for lifelong supplements and follow-up. The procedure choice was left up to the patient after a detailed formal presentation of the anatomy, mechanisms of action, short- and long-term complication rates and expected weight loss associated with each procedure.

Patients with American Society of Anaesthesiology (ASA) class 4 disease or an obesity surgery mortality risk score (OS-MRS¹⁷) greater than 4 were excluded from having surgery at the SHCC. Patients weighing more than 205 kg were also excluded owing to ambulance transfer restrictions. The surgeon was the only constant at the THCC, with patients treated on the ward or in the ICU as deemed appropriate. All patients were placed on a high-protein, low-carbohydrate diet for 2 weeks before surgery. Thromboprophylaxis included sequential compression stockings during the surgery, the administration of 5000 units of unfractionated heparin subcutaneously every 12 hours until discharge, and very aggressive and early mobilization, especially at the SHCC, where the patient-to-staff ratio made this feasible. No perioperative antibiotics were given. We followed "awake intubation" protocol for all patients at the SHCC using fiberoptic intubation under heavy sedation to secure the airway before administration of paralytic agents.

Our laparoscopic gastric bypass technique involves a 30–50 cm biliopancreatic limb and a 100 cm retrocolic, antegastric, alimentary limb. The jejunojunal anastomosis is constructed side-to-side with a single firing of a linear endostapler, and the stapler entry point defect is hand sewn. The gastric pouch is small (2 × 7 cm) and vertically oriented, and the gastrojejunal anastomosis is hand sewn. Pneumatic and methylene blue tests are used to ensure the integrity of staple and suture lines at the time of surgery. The Petersen space and the transverse mesocolic defect are routinely closed with running polypropylene sutures. Vertical sleeve gastrectomy is performed "loosely" over a 36 F bougie using the appropriate endostapler with no reinforcements. Integrity of the staple line is verified, as for

the gastric bypass. All gastric bands are inserted using the pars flaccida technique.

The outcome data (demographic characteristics, weight loss, complications) were recorded prospectively in our bariatric surgery registry. We calculated the kilogram weight loss, percent body weight loss (weight loss ÷ baseline body weight × 100) and percent excess weight loss (%EWL; $100\% \times [(W_0 - W_i) \div EW_0]$, where W_0 is the weight in kilograms at the time of surgery, W_i is the weight in kilograms at the last follow-up, and EW_0 is the excess weight at the time of surgery). Excess weight was estimated according to the formula described by Deitel and Greenstein¹⁸ and is based on the Metropolitan Tables for middle frame individuals. Complications occurring within 30 days of the date of surgery were defined as short-term complications, and complications occurring after 30 days were defined as long-term. We also determined 30-day mortality and the number of deaths during the long-term follow-up that could be related to the original bariatric surgery.

Statistical analysis

We used IBM SPSS Statistics 20 for the computations and statistical analysis. Continuous variables were tested for significance using unpaired *t* tests; the χ^2 was used to compare proportions. Logistic regression analysis was used to compare individual variable contribution to mortality.

RESULTS

In all, 830 patients were enrolled in the study: 676 (81.4%) were treated at the SHCC and 154 (18.6%) were treated at the THCC. Table 1 shows the patient demographic and clinical characteristics and the type of surgery performed at each centre. There were significantly more women treated at the SHCC than the THCC, and patients treated at the SHCC were slightly younger by

about 1 year than those treated at the THCC. The patients at the THCC were heavier and their BMI significantly higher than those treated at the SHCC. Laparoscopic gastric bypass was the most common procedure (85.4%), followed by gastric banding and gastric sleeve. There was a significant discrepancy in the distribution of the surgeries performed at each centre; gastric bypass was the predominant procedure because gastric banding was not publicly funded in Quebec until recently.

The distribution of obesity-associated comorbidities is shown in Table 2. Most comorbidities were equally distributed; however, heart disease was more common among patients treated at the THCC, as was expected based on patient selection criteria, and weight-bearing osteoarthritis was also more common among those treated at the THCC, as was expected based on the higher BMI in that group. Table 3 shows the types of complications recorded within the first 30 days after the surgery. As expected, complication rates were slightly higher in the THCC group given the patient selection criteria. Obesity surgery mortality risk scores and ASA score were also significantly higher in the THCC group (Table 4).

The major complication rate was 2.3% at the SHCC and 5.8% at the THCC ($p = 0.036$). Similarly, minor complications were significantly more frequent in the THCC group ($p = 0.003$). The readmission rates in the 30 days after surgery were equivalent between the groups. There were 7 direct transfers (1.2%) from the SHCC to the THCC via the dedicated “service corridor.” The reasons for the transfer as well as the treatment and outcomes are shown in Table 5. All 7 patients made a full recovery. An additional 9 patients returned to the hospital on average 11 (standard deviation [SD] 9, range 3–33) days after surgery for the reasons outlined in Table 6. All of them were successfully treated and made a full recovery.

Table 1. Demographic and clinical characteristics of the study population

Factor	Group; no. (%) or mean ± SD (range)*			p value
	Total (%) n = 830	SHCC n = 676	THCC n = 154	
Sex				0.003
Men	260 (31.3)	197 (29.1)	63 (40.9)	
Women	570 (68.7)	479 (70.9)	91 (59.1)	
Gastric bypass	709 (85.4)	558 (82.5)	151 (98.1)	0.001
Gastric band	92 (11.1)	89 (13.2)	3 (3.3)	0.001
Gastric sleeve	29 (3.5)	29 (4.3)	0	0.001
Age, yr	43.3 ± 11.4 (13–79)	42.1 ± 10.1 (13–79)	43.6 ± 11.8 (17–66)	0.031
Weight, kg	138.0 ± 28.9 (49.9–245.9)	133.8 ± 25.3 (49.5–218.4)	156.9 ± 35.6 (96.4–245.9)	0.001
Body mass index	48.8 ± 8.3 (19.4–84.8)	47.5 ± 7.4 (19.4–71.9)	54.4 ± 9.7 (36.7–84.8)	0.001

SD = standard deviation; SHCC = secondary health care centre; THCC = tertiary health care centre.
*Unless otherwise indicated.

Table 2. Obesity-associated diseases and conditions in the 2 cohorts

Disease/condition	Group, no. (%)		p value
	SHCC	THCC	
Asthma	141 (21)	39 (26)	NS
Chronic back pain	545 (81)	128 (85)	NS
Depression	278 (41)	64 (43)	NS
Fatigue	377 (56)	95 (63)	NS
Heart disease, any type	128 (19)	42 (27.8)	0.012
High triglycerides	85 (12.6)	14 (9.3)	NS
Hypertension	275 (41)	56 (37)	NS
Osteoarthritis of joints	292 (43)	91 (60)	0.001
Short of breath on exertion	419 (62.2)	99 (65.6)	NS
Sleep apnea on CPAP	56 (8.3)	14 (9.3)	NS
Stress incontinence	235 (35)	61 (40.4)	NS
Type 2 diabetes	144 (21)	31 (21)	NS

CPAP = continuous positive airway pressure; NS = not significant; SHCC = secondary health care centre; THCC = tertiary health care centre.

There were 2 deaths at the THCC. The first was a woman with a BMI of 75.5, ASA of 3 and OS-MRS of 3 who underwent an uneventful laparoscopic gastric bypass but suffered a massive pulmonary embolism on day 4 post-surgery, the day of her planned discharge. The second was a woman with a BMI of 59, ASA of 4 and OS-MRS of 4, who was on home oxygen and who underwent an uneventful laparoscopic gastric bypass. She was admitted to the ICU as planned for progressive weaning from the ventilator, but nosocomial pneumonia developed and she died in the ICU. No deaths occurred in the SHCC. Overall mortality was 0.2%. Logistic regression analysis failed to identify any variables (age, sex, location of surgery, starting BMI, ASA score, OS-MRS) contributing to the risk of death owing to low incidence of death.

Weight loss in kilograms (Fig. 1) and the percentage of total weight loss (Fig. 2) were equivalent between the 2 centres. This method of weight loss data presentation is not affected by the starting patient weight or BMI. The

%EWL (Fig. 3) appears to be better at the SHCC than the THCC. Since the %EWL is affected by the starting BMI (i.e., better %EWL in patients with lower BMI), we adjusted the %EWL calculation for starting BMI and found no difference between the 2 groups.

Figure 4 shows a comparison of the excess weight loss between the 3 laparoscopic bariatric surgery procedures at the SHCC (there were insufficient data to compare the 3 procedures at the THCC). Gastric bypass was associated with significantly better weight loss results than the gastric band and gastric sleeve procedures.

DISCUSSION

The number of Canadians who are morbidly obese is estimated as 3% of 34 million (2010 data from Statistics Canada) or about 1 million people. If 10% of morbidly obese people in Canada were to request bariatric surgery to treat obesity in the next 10 years, this would mean performing about 10 000 operations per year. However, this estimate does not take into account the exponential increases in the prevalence of obesity and type 2 diabetes that have been projected for the coming 10–20 years.¹⁹

A recent survey of centres known to perform more than 20 bariatric surgical procedures per year conducted by the MUHC team provides the best available data on the volume of bariatric surgery in Canada.¹² The survey results showed that the 12 Canadian centres that responded performed 1313 procedures in 2007. Given that 6783 patients were waiting for bariatric surgery, the authors estimated the wait time for bariatric surgery in Canada to be 5.2 years.

The Ontario Ministry of Health and Long-Term Care stated in their December 2005 report that the province should increase its bariatric surgery capacity to at least 3500 surgeries per year, as did the AETMIS recommendation. This latter report, released in 2005, recommended

Table 3. The types of complications recorded within the first 30 days after surgery at each site

Complication	Group; no. (%)	
	SHCC, n = 676	THCC, n = 154
Major complications	16 (2.3)	9 (5.8)
Death	0	2 (1.3)
Anastomotic leak	10 (1.4)	4 (2.5)
SIRS response, no leak detected at laparoscopy	2 (0.3)	1 (0.6)
Abdominal abscess	1 (0.1)	2 (1.3)
Colon perforation	1 (0.1)	0
Jejunojunostomy leak	1 (0.1)	0
Acute gastric dilatation	1 (0.1)	0
Minor complications	35 (5.2)	19 (12.3)
Stenosis of the gastrojejunostomy	5 (0.7)	2 (1.3)
Abdominal wall hematoma	3 (0.4)	2 (1.3)
Bleeding from a staple line	3 (0.4)	1 (0.6)
Liver laceration (minor)	4 (0.6)	3 (1.9)
Fever NYD	4 (0.6)	0
Port site infection	4 (0.6)	0
Pneumonia	2 (0.3)	2 (1.3)
Pulmonary edema	1 (0.1)	3 (1.9)
Small bowel obstruction	2 (0.3)	1 (0.6)
Perisplenic hematoma	1 (0.1)	2 (1.3)
Unable close mesenteric defects	1 (0.1)	2 (1.3)
Pneumatic test positive, repaired in OR	1 (0.1)	1 (0.6)
Renal calculus, hematuria	1 (0.1)	1 (0.6)
Acute diverticulitis	0	1 (0.6)
Advanced cirrhosis of liver found at surgery	1 (0.1)	0
Camera port infection	1 (0.1)	0
Neurapraxia left arm	1 (0.1)	0
Pancolitis	1 (0.1)	0
Ruptured alimentary limb by MB test	1 (0.1)	0
Allergic reaction to unknown trigger	1 (0.1)	0
Staple malfunction	0	1 (0.6)

MB = methylene blue; NYD = not yet diagnosed; OR = operating room; SHCC = secondary health care centre; SIRS = systemic inflammatory response syndrome; THCC = tertiary health care centre.

Table 4. Patient stratification and complications recorded within the first 30 days after surgery at each site

Factor	Group; mean ± SD or no. (%)*		
	SHCC	THCC	p value
OS-MRS	1.6 ± 1.1	2.6 ± 1.7	0.001
ASA class	2.8 ± 0.8	3.2 ± 0.8	0.001
Operating time in-out of room, min.	89.0 ± 12.0	145.0 ± 23.0	0.002
Length of stay, d	1.9 ± 0.1	2.8 ± 0.4	0.003
Major complications	16 (2.3)	9 (5.8)	0.036
Minor complications	35 (5.2)	19 (12.3)	0.003
Readmission within 30 d of surgery	16 (2.3)	3 (1.9)	NS
Direct transfers to THCC	7 (1.2)	—	—
Deaths	0	2 (1.3)	NS

ASA = American Society of Anaesthesiologists; OS-MRS = obesity surgery mortality risk score; SHCC = secondary health care centre; THCC = tertiary health care centre.
*Unless otherwise indicated.

that that the province of Quebec substantially increase its capacity for bariatric surgery from 716 operations per year in 2005 to more than 3500 by 2012 to meet demand.

Since the release of these reports, Ontario has increased bariatric surgery capacity in the public system to an estimated 2085 per year in 2011, and Quebec has increased capacity to an estimated 1757 per year. Both estimates are below the targets. The present study was initiated to determine if one of the AETMIS report recommendations could be implemented: the strategy of designating several SHCCs and mandating them to complete about 200 select cases of bariatric surgery each year. Each centre would be linked by a “service corridor” to a THCC within each of the 4 integrated health networks. With the addition of 10

such SHCCs (200 cases each) to the 4 THCCs (500 cases each) within Quebec’s 4 integrated health networks, the yearly bariatric surgery capacity could be increased to 4000. The SHCCs could include some specialized medical centres¹⁵ that deliver publicly funded, private delivery bariatric surgeries according to law 33 enacted in 2006. We have now collected sufficient statistics to suggest that, with proper patient selection, this approach could be feasible.

Patient selection criteria allow for safe surgery to be delivered at SHCCs with acceptable mortality and short-term and long-term complications. Overall mortality and centre-specific mortality are well within accepted values.²⁰ Weight outcomes are also favourable. Weight loss greater than 50% of the excess weight²¹ and reduction of the BMI²²

Table 5. Reason for direct transfer from the SHCC to the THCC

No.	Year	BMI	Sex	Surgery	Day of transfer	Reason for transfer
1	2006	40	M	Gastric bypass	2	Latrogenic colon perforation by endoshear at time of surgery. Systemic inflammatory response developed on postoperative day 2. Required partial colon resection and ICU stay.
2	2007	47	M	Gastric bypass	1	Systemic inflammatory response. No leak visible on CT scan, but perisplenic hematoma present. Staple line leak developed on postoperative day 28. Treated with drainage and stent.
3	2008	43	F	Gastric bypass	1	Jejunojejunostomy rotation and obstruction with pouch dilation and staple line leak. Laparotomy and repair of both.
4	2009	60	F	Gastric sleeve	2	Systemic inflammatory response. No leak visible on CT scan. Uneventful discharge.
5	2010	51	M	Gastric bypass	1	Systemic inflammatory response on postoperative day 2 due to an ischemic perforation of the gastric pouch. This was a revision surgery: band to bypass. Laparoscopic suture repair.
6	2010	46	M	Gastric bypass	1	Ventricular fibrillation developed immediately upon reversal of anesthesia while the patient was still intubated. Defibrillated successfully and transferred to THCC for full cardiac workup, including cardiac angiography. No pathology found. Complete recovery.
7	2011	47	M	Gastric bypass	1	Systemic inflammatory response developed on postoperative day 1. Leak or other cause not visible on CT scan or during laparoscopy. Discharged on postoperative day 5. Uneventful recovery.

BMI = body mass index; CT = computed tomography; ICU = intensive care unit; SHCC = secondary health care centre; THCC = tertiary health care centre.

Table 6. Reason for readmission to the THCC after discharge from the SHCC

No.	Year	BMI	Sex	Surgery	Day post discharge	Reason for readmission
1	2006	46	F	Gastric bypass	33	Perforated jejunojejunostomy that required open repair.
2	2007	53	F	Gastric bypass	15	Staple line leak requiring laparoscopic repair.
3	2007	57	F	Gastric bypass	6	Subcutaneous abscess at operating port site. Drained through port site.
4	2008	37	F	Gastric bypass	3	Severe systemic inflammatory response. No leak demonstrated on CT scan or at laparoscopy. Discharged 2 days later.
5	2008	41	M	Gastric bypass	6	Moderate systemic inflammatory response. No leak visible on CT scan, but a suspected “infected hematoma” was drained. No other evidence of leak. Discharged in 1 week.
6	2008	51	F	Gastric sleeve	7	Staple line leak, upper end. Computed tomography-guided drainage.
7	2009	42	M	Gastric bypass	10	Moderate systemic inflammatory response. Suspected subphrenic abscess on CT scan. Laparoscopy failed to identify the leak. Full recovery.
8	2009	38	F	Gastric bypass	7	Multiple previous surgeries. Acute bowel obstruction developed distal to the jejunojejunostomy. Disruption of anastomosis. Laparotomy and repair.
9	2011	41	F	Gastric sleeve	12	Multiple previous surgeries. Enterotomy during laparoscopic adhesion dissection. Acute bowel obstruction developed postsurgery and disrupted enterotomy repair. Laparotomy and repair.

BMI = body mass index; CT = computed tomography; F = female; M = male; SHCC = secondary health care centre; THCC = tertiary health care centre.

to less than 35 have been proposed as potential definitions of success of a bariatric surgical procedure. The medical obesity literature uses actual weight loss and percent body weight loss. The weight loss results from the present study are reported using all these outcome measures and show that the weight loss is robust and maintained for the 4 years of follow-up available in the SHCC. These results suggest superior weight loss after gastric bypass than gastric band or sleeve.

Limitations

This study has several limitations. It represents the personal series of 1 experienced bariatric surgeon's minimally

invasive laparoscopic bariatric surgery practice, which is well beyond the learning curve for bariatric procedures.²³ As such, there is no surgeon- or technique-related variability. Less experienced surgeons may not be able to duplicate these results. This is why it is imperative to select the SHCC very carefully and provide adequate training to the preoperative, perioperative and postoperative teams and surgeons who will be caring for these patients. This is not a randomized study and, as such, it is subject to all the potential bias of a prospective cohort study. The perioperative follow-up was adequate, and no deaths or life-threatening complications could have been missed. Despite determined efforts of long-term follow-up for all patients, this was not successful. Finally, resolution




Fig. 1. Weight loss in kilograms in patients with laparoscopic gastric bypass at each site. Mean and standard deviation are shown as well as the number of cases followed-up at each time point. SHCC = secondary health care centre; THCC = tertiary health care centre.




Fig. 3. Percent excess weight loss in patients with laparoscopic gastric bypass at each site. Mean and standard deviation are shown as well as the number of cases followed-up at each time point. SHCC = secondary health care centre; THCC = tertiary health care centre.




Fig. 2. Percent total body weight loss in patients with laparoscopic gastric bypass at each site. Mean and standard deviation are shown as well as the number of cases followed-up at each time point. SHCC = secondary health care centre; THCC = tertiary health care centre.




Fig. 4. Comparison of the excess weight loss between the 3 laparoscopic bariatric surgery procedure sat the secondary health centre (insufficient data to compare the tertiary health care centre. Mean and standard deviation are shown.

and improvement of comorbidities were not included in the analysis, as the aim of this study was to concentrate on weight loss and morbidity and mortality assessment.

CONCLUSION

With proper patient selection, a well-trained, dedicated health care team and a “service corridor” to an affiliated THCC, laparoscopic bariatric surgery, including gastric bypass, can be performed safely in an SHCC. This service model warrants more study to determine whether it can be widely applied across Canada.

Competing interests: None declared.

References

1. Torrance GM, Hooper MD, Reeder BA. Trends in overweight and obesity among adults in Canada (1970–1992): evidence from national surveys using measured height and weight. *Int J Obes Relat Metab Disord* 2002;26:797-804.
2. Katzmarzyk PT. The Canadian obesity epidemic, 1985–1998. *CMAJ* 2002;166:1039-40.
3. Mokdad AH, Marks JS, Stroup DF, et al. Actual causes of death in the United States, 2000. *JAMA* 2004;291:1238-45.
4. Christou NV, Look D, Maclean LD. Weight gain after short- and long-limb gastric bypass in patients followed for longer than 10 years. *Ann Surg* 2006;244:734-40.
5. Brolin RE. Bariatric surgery and long-term control of morbid obesity. *JAMA* 2002;288:2793-6.
6. Christou NV, Sampalis JS, Liberman M, et al. Surgery decreases long-term mortality, morbidity, and health care use in morbidly obese patients. *Ann Surg* 2004;240:416-23; discussion 423-44.
7. Busetto L, Mirabelli D, Petroni ML, et al. Comparative long-term mortality after laparoscopic adjustable gastric banding versus nonsurgical controls. *Surg Obes Relat Dis* 2007;3:496-502, discussion 502.
8. Peeters A, O'Brien PE, Laurie C, et al. Substantial intentional weight loss and mortality in the severely obese. *Ann Surg* 2007;246:1028-33.
9. Sjöström L, Narbro K, Sjöström CD, et al. Effects of bariatric surgery on mortality in Swedish obese subjects. *N Engl J Med* 2007;357:741-52.
10. Adams TD, Gress RE, Smith SC, et al. Long-term mortality after gastric bypass surgery. *N Engl J Med* 2007;357:753-61.
11. Sampalis JS, Liberman M, Auger S, et al. The impact of weight reduction surgery on health-care costs in morbidly obese patients. *Obes Surg* 2004;14:939-47.
12. Christou NV, Efthimiou E. Bariatric surgery waiting times in Canada. *Can J Surg* 2009;52:229-34.
13. Padwal RS, Sharma AM. Treating severe obesity: morbid weights and morbid waits. *CMAJ* 2009;181:777-8.
14. Surgical treatment of morbid obesity — an update. Agence d'évaluation des technologies et des modes d'intervention en santé. Available: www.iness.qc.ca/fileadmin/doc/AETMIS/Rapports/Obesite/2005_04_en.pdf (accessed 2013 July 3).
15. Centre médical spécialisé. In: website of Santé et services sociaux Québec. Available: www.msss.gouv.qc.ca/sujets/organisation/cms/index.php?centre_medical_specialise (accessed 2012 Feb. 29).
16. Grundy SM, Barondess JA, Bellegie NJ, et al.; Consensus Development Panel. Bethesda (MD): Office of Disease Prevention; 1991. *Gastrointestinal surgery for severe obesity: National Institutes of Health Consensus Development Conference Statement*. Available: <http://consensus.nih.gov/1991/1991GISurgeryObesity084html.htm> (accessed 2013 Mar. 22).
17. DeMaria EJ, Portenier D, Wolfe L. Obesity surgery mortality risk score: proposal for a clinically useful score to predict mortality risk in patients undergoing gastric bypass. *Surg Obes Relat Dis* 2007;3:134-40.
18. Deitel M, Greenstein RJ. Recommendations for reporting weight loss. *Obes Surg* 2003;13:159-60.
19. Somerville R. *An economic tsunami: the cost of diabetes in Canada*. Toronto (ON): Canadian Diabetes Association; 2009. Available: www.diabetes.ca/documents/get-involved/FINAL_Economic_Report.pdf (accessed 2010 Dec. 21).
20. Nguyen NT, Masoomi H, Laugenour K, et al. Predictive factors of mortality in bariatric surgery: data from the Nationwide Inpatient Sample. *Surgery* 2011;150:347-51.
21. Reinhold RB. Critical analysis of long term weight loss following gastric bypass. *Surg Gynecol Obstet* 1982;155:385-94.
22. Biron S, Hould FS, Lebel S, et al. Twenty years of biliopancreatic diversion: What is the goal of the surgery? *Obes Surg* 2004;14:160-4.
23. Andrew CG, Hanna W, Look D, et al. Early results after laparoscopic Roux-en-Y gastric bypass: effect of the learning curve. *Can J Surg* 2006;49:417-21.