

PERIOPERATIVE MORTALITY AFTER PNEUMONECTOMY: ANALYSIS OF RISK FACTORS AND REVIEW OF THE LITERATURE

Daniel E. Swartz, MD; Kevin Lachapelle, MD; John Sampalis, PhD; David S. Mulder, MD; Ray C.-J. Chiu, MD, PhD; James Wilson, MD

OBJECTIVE: To determine risk factors for perioperative death associated with pneumonectomy.

DESIGN: A retrospective case-control study in which a perioperative death group was compared with a survivor group, and a review of the English literature on the subject.

SETTING: The Montreal General Hospital, a tertiary-care teaching institution.

PATIENTS AND INTERVENTION: Ninety-two consecutive patients who underwent pneumonectomy between April 1989 and 1994.

MAIN OUTCOME MEASURES: The effects of age, sex, smoking history, tumour size, type and stage, pulmonary function, cardiovascular risks, comorbidity, preoperative blood values and volume of fluids administered perioperatively. Values from the literature were reported for comparison.

RESULTS: The perioperative death rate was 10.9%. Selection bias and in-hospital values reported in the literature have underestimated the death rate, with actual rates ranging from 7% to 11%. Age (odds ratio 2.48, $p = 0.04$), the presence of 1 or more comorbid diseases (odds ratio 7.92, $p = 0.05$) and amount of fluids given in the first 12 hours postoperatively (odds ratio 2.21, $p = 0.06$) were found to be significant risk factors for death. Multivariate logistic regression demonstrated that the volume of fluids given remains an independent risk factor whereas age and comorbid disease are dependent variables.

CONCLUSIONS: The results were consistent with previously reported death rates and risk factors. Patient age and concomitant disease are not modifiable risk factors, but perioperative fluid administration and other means to prevent postpneumonectomy pulmonary edema may reduce the perioperative death rate.

OBJECTIF : Déterminer les facteurs de risque de mortalité périopératoire liée à la pneumonectomie.

CONCEPTION : Étude cas-témoin rétrospective au cours de laquelle on a comparé un groupe de patients morts en période périopératoire à un groupe de survivants, et recension des écrits en anglais sur la question.

CONTEXTE : Hôpital Général de Montréal, établissement d'enseignement de soins tertiaires.

PATIENTS ET INTERVENTIONS : Quatre-vingt-douze patients consécutifs qui ont subi une pneumonectomie entre avril 1989 et 1994.

PRINCIPALES MESURES DES RÉSULTATS : Effets de l'âge, sexe, antécédents de tabagisme, grosseur, type et stade de la tumeur, fonction pulmonaire, risques cardiovasculaires, comorbidité, valeurs sanguines préopératoires et volumes de liquides administrés en période périopératoire. On a indiqué les valeurs tirées des écrits pour des fins de comparaison.

RÉSULTATS : Le taux de mortalité périopératoire s'est établi à 10,9 %. Le biais de la sélection et les valeurs à l'hôpital indiquées dans les écrits entraînent une sous-estimation des taux de mortalité, les taux réels variant de 7 % à 11 %. On a constaté que l'âge (coefficient de probabilité de 2,48, $p = 0,04$), la présence d'une ou plusieurs maladies entraînant un état comorbide (coefficient de probabilité de 7,92, $p = 0,05$) et les vo-

From the Department of Surgery, Montreal General Hospital and McGill University, Montreal, Que.

Accepted for publication Feb. 13, 1997

Correspondence to: Dr. Ray C.-J. Chiu, Director of Cardiovascular and Thoracic Surgery, Montreal General Hospital, 1650 Cedar Ave., Montreal QC H3G 1A4; tel 514 934-2839; fax 514 934-8289

© 1997 Canadian Medical Association (text and abstract/résumé)

lumes de liquides administrés au cours des 12 heures qui ont suivi l'intervention (coefficient de probabilité de 2,21, $p = 0,06$) constituaient d'importants facteurs de risque de décès. Une régression logistique multivariée a démontré que les volumes de liquides administrés demeurent un facteur de risque indépendant tandis que l'âge et la comorbidité sont des variables dépendantes.

CONCLUSIONS : Les résultats ont été conformes aux taux de mortalité et aux facteurs de risque signalés auparavant. L'âge du patient et la présence d'une comorbidité sont des facteurs de risque non modifiables, mais l'administration de liquides pendant la période périopératoire et d'autres moyens de prévenir l'œdème pulmonaire consécutif à une pneumonectomie peuvent réduire le taux de mortalité périopératoire.

Pneumonectomy has among the highest morbidity and mortality of thoracic procedures. The perioperative death rate, defined by The Society of Thoracic Surgeons as death within 30 days of operation, has been reported as high as 36% in the recent scientific literature.¹ Although studies have documented lower death rates,²⁻⁵ a closer review of the literature reveals that a selection bias of patients for study exists and contributes to falsely low death rates. Many preoperative parameters have been investigated for their contribution to perioperative death with disappointing consistency, and at present no general consensus exists for patient selection. This is compounded by the fact that the decision to perform a pneumonectomy is sometimes made intraoperatively, which obviates the use of selection criteria. However, one can usually determine preoperatively that a pneumonectomy may be necessary and excessive risk might persuade the surgeon to pursue nonoperative treatments. The aim of this paper is to review the literature on pneumonectomy over the past 25 years with respect to study designs and outcomes to identify parameters that present a significant risk of perioperative death after pneumonectomy. We reviewed our experience with pneumonectomy between 1989 and 1994 and included the data in this review.

HISTORICAL OVERVIEW

The pneumonectomy procedure has survived with little modification since its initial troublesome beginnings. Harvey, Erdman and Beattie⁶ in

a recent review of the technique provided an extensive history of pneumonectomy. Reinhoff in 1933 performed a pneumonectomy using individual ligation of hilar structures and covered the bronchus with a flap of parietal pleura.⁷ No drainage or thoracoplasty was done. During this time, Shenstone in Toronto described a similar technique,⁸ which has remained essentially unchanged. Pneumonectomy was believed to be the standard curative resection for lung cancer through the 1940s. Equal cure rates were subsequently demonstrated from more limited resections with improved morbidity and mortality. Lobectomy is now the curative procedure of choice for resectable lung cancer whereas pneumonectomy is indicated when lesser resections are not possible.

The perioperative death rate has decreased substantially since the early days of pneumonectomy when survival was the exception. Wilkins, Scannell and Craver⁷ reviewed the Massachusetts General Hospital data of 392 pneumonectomies performed between 1931 and 1970 and demonstrated in-hospital death rates of 56.5% for the decade 1931 to 1940; 14.7% between 1941 and 1950; 13.8% between 1951 and 1960; and 11.1% for the decade 1961 to 1970. This trend approximates the death rates from previous decades. Of note is the authors' use of "in-hospital" deaths to estimate perioperative (30-day) mortality. Many retrospective analyses use in-hospital data,^{2,3,7-11} since many patients are discharged before 30 days and hospital charts hold no subsequent record of patient status. In their analysis of hos-

pital-associated deaths calculated from discharge data, Jencks, Williams and Kay¹² determined that in-hospital mortality differs significantly from 30-day mortality because of bias in length of stay. Because many pneumonectomy patients are discharged before 30 days, in-hospital mortality underestimates perioperative deaths.

LITERATURE SURVEY

A summary of the English language literature on perioperative death rates associated with pneumonectomy over the past 20 years appears in Table I.^{1-5,7-11,13-20} A MEDLINE search was used to find papers indexed with the subject "pneumonectomy" and limited by the heading "mortality." Papers in foreign languages and in journals not available in the city of Montreal were included if the abstracts were printed in English and reported death rates.

A wide range of perioperative death rates have been reported (5% to 23%). However, closer inspection revealed the presence of both selection and observation biases. We have previously discussed the observation bias related to in-hospital versus 30-day death rates. Ginsberg and associates⁴ reported the often-quoted 30-day death rates for the 2220 resections in patients enrolled in the 5 centers and 12 institutions of the Lung Cancer Study Group. Of the 569 pneumonectomies performed during a 3-year period among 12 institutions with death rates ranging from 0% to 25%, the authors noted an average death rate of 6.2%. This study and others^{1,2,5,8,13-16} demonstrated selection bias by limiting crite-

ria for entry into the study series to patients with primary lung cancer. When analyses of perioperative death rates for pneumonectomy included all pneumonectomies rather than patients with selected diseases, the death rate was usually higher (Table I) as was demonstrated in 2 published series^{5,17} from the same institution over the same time period. One study was limited to patients presenting with primary bronchogenic carcinoma; the other included all comers. The perioperative mortalities for the 2 groups were 5% and 7%, respectively. The prevailing view that pneumonectomy for benign disease carries a higher death rate than for malignant disease has led to the separation of these 2 categories of patients in most studies. However, a recent retrospective study by Conlan and associates³ noted a

1.6% death rate for 124 consecutive pneumonectomies for benign disease. Others have included both benign and malignant diseases in their studies.^{9,18,21} Subclassifications of patients in previous studies (i.e., benign v. malignant, high risk v. low risk, age over v. under 70 years) are noted where indicated in Table I.^{1-5,7-11,13-20} Acceptable 30-day death rates after pneumonectomy as reported in the literature over the last 7 years, whether for primary cancer or all diseases, have ranged from 7% to 11%. Completion pneumonectomies were not included in this review.

We recently reviewed all pneumonectomies at the Montreal General Hospital over the 5 years from April 1989 to 1994. The goal of this study was to determine risk factors for perioperative death.

METHODS

The study design was similar to that of many other analyses of perioperative death after pneumonectomy.^{5,6,10,11,17,18} The patients were assigned to either “survivor” or “death” groups based on their status 30 days after operation. Table II lists the parameters that were investigated for significance as risk factors for perioperative death. The means (and standard deviations) of continuous data were compared between groups using a *t*-test. Then clinically acceptable cut-off values were applied to convert the data into categorical values before statistical significance was assessed by χ^2 analysis. Parameters that were significantly different for either or both tests were considered significant risk factors for perioperative death. Univariate and multi-

Table I

Reported Perioperative Deaths After Pneumonectomy

Series	No. of patients	Duration of study, yr	Death rate, %	Comments
Weiss, 1974 ¹³	212	5	17	Males, primary cancer only
Kirsh et al, 1976 ¹⁹	6	10	17	Primary and metastatic cancers
Wilkins, Scannell and Craver, 1978 ⁷	77	7	9.1	In-hospital, primary lung cancer
Nagasaki, Flehinger and Martini, 1982 ¹⁴	72	8	7	Primary lung cancer only
Ginsberg et al, 1983 ⁴	569	3	6.2	Primary lung cancer only
Keagy et al, 1983 ¹⁵	90	10	12.2	Primary lung cancer only
Kohman et al, 1986 ⁸	94	12.5	11.7	In-hospital, primary lung cancer,
Krowka et al, 1987 ⁹	244	5	11	Benign, malignant, includes completion, in-hospital
Wahi et al, 1989 ¹⁷	197	5	7	Primary and metastatic lung cancers
Putnam et al, 1990 ⁵	139	5	5	Primary lung cancer only
Roxburgh, Thompson and Goldstraw, 1991 ²	105	4	6.7	In-hospital, primary lung cancer
Whittle et al, 1991 ²⁰	—	2	17	Age > 69 yr, 30-d, cancer, in-hospital, primary lung cancer
Patel, Townsend and Fountain, 1992 ¹⁰	197	5	8.6	In-hospital, primary lung cancer
Romano and Mark, 1992 ¹¹	1529	3	11.7	In-hospital, primary lung cancer
Pierce et al, 1994 ¹	11	1.5	36.4	Primary lung cancer
Conlan et al, 1995 ³	124	15	3	In-hospital, benign only, includes completion
Ferguson, Reeder and Mick, 1995 ¹⁸	92	13	12	Benign and malignant
Shah et al, 1996 ¹⁶	59	2	1.7	Stages I and II primary cancer only

variate logistic regression analyses were performed to determine incremental risk as independent variables and adjusted for other risk factors.

RESULTS

After excluding pneumonectomy done for organ harvest and lung transplantation, we were left with 92 pneumonectomies performed for the following indications: primary bronchogenic carcinoma (87, of which 83 were non-small cell, 2 small cell and 2 carcinoids), mesothelioma (1), metastatic carcinoma (1), aspergilloma (1), cystic fibrosis (1) and emphysema with bronchopleural fistula (1). There were 10 deaths for a 30-day perioperative death

rate of 11% (Table III). Of 2 patients operated on for benign diseases there was 1 death, whereas of 90 patients operated for lung cancer there were 9 deaths. With respect to the characteristics of the 2 groups (Table IV), age and the presence of 1 or more comorbid states were significantly greater in the death group. Significant risk factors for perioperative death were: advancing age, volume of fluids administered in the first 12 hours postoperatively and the presence of 1 or more comorbid states (Table IV), which included coronary artery disease, arrhythmias, congestive heart failure, hypertension, peripheral vascular disease, diabetes, chronic obstructive pulmonary disease, restrictive lung disease, extrapulmonary can-

cer, chronic liver disease, chronic renal failure or inflammatory bowel disease. Univariate and multivariate logistic regression analyses for the 3 factors are presented in Table V.

DISCUSSION

Five of the 10 patients who died had autopsies performed. Two patients suffered cardiac herniation through pericardial defects that led to a refractory arrhythmia in 1 and a myocardial infarction in the other; autopsy in the latter revealed diffuse infarction and severe (75% to 99%) stenosis of the 3 coronary arteries by atheromatous plaques (Table III). Three patients had postpneumonectomy pulmonary edema (PPE) recorded as the cause of death, and 3 others died secondary to adult respiratory distress syndrome (ARDS) and multisystem organ failure (MSOF) with or without sepsis. Because hard diagnostic criteria may be lacking, PPE is an often unrecognized but fatal complication of pneumonectomy. It is possible that the last 3 patients had PPE and secondarily ARDS, sepsis and MSOF.

Other investigators have studied the aforementioned risk factors for perioperative death after pneumonectomy. However, the results have lacked consistency. Not for lack of analysis does it remain unclear whether advancing age carries a significant risk for perioperative death. Several authors compared patients older and younger than 70 years and found no significant difference with respect to death between the 2 groups,^{2,4,10,22} whereas others demonstrated a significant difference in the death rate for older patients.^{11,14,18} Our data, reported in Table IV, demonstrated statistical significance for age as a risk factor when examined on a continuous spectrum (survivors 60.8 [11.2] years v. death

Table II

Parameters Assessed for Risk of Death After Pneumonectomy

Age
Smoking history (pack-years, years stopped smoking before operation)
Significant weight loss
Primary diagnosis (non-small cell bronchogenic cancer, metastatic disease, inflammatory disease)
Side of resection
Size of tumour
Cell type of tumour (if applicable)
Stage (clinical and pathologic)
Pulmonary function tests (FEV ₁ , % predicted FEV ₁ , FVC, % predicted FVC, FEV ₁ /FVC, TLC, RV, D _{LCO})
Number of cardiac risk factors (smoking, diabetes, hypertension, hyperlipidemia, history of cardiac disease)
Significant medical history (cardiac, diabetes, hypertension, respiratory, pulmonary cancer, peripheral vascular disease, liver disease, renal insufficiency, inflammatory bowel disease)
Preoperative serum measurements (hemoglobin, albumin, alkaline phosphatase)
Abnormal electrocardiogram
Preoperative arterial blood gas levels (pH, P _{CO₂} , P _{O₂} , bicarbonate, % oxygen saturation)
Anesthetic time
Estimated blood loss
Blood transfusion
Postoperative fluid administration (0–12, 12–24, 24–48 and 48–72 h)
Postoperative urine output (0–12, 12–24, 24–48 and 48–72 h)
Number of days in-hospital

FEV₁ = forced expiratory volume in the first second, FVC = forced vital capacity, TLC = total lung capacity, RV = residual volume, D_{LCO} = diffusing capacity of the lung for carbon monoxide, P_{CO₂} = partial pressure of carbon dioxide, P_{O₂} = partial pressure of oxygen

68.6 [8.2] years, $p = 0.04$), categorically for patients younger than 70 years versus those 70 years of age or older (odds ratio [OR] 6.2, 95% confidence interval [CI] 1.6 to 24.5, $p = 0.005$), and in a univariate logistic regression analysis with yearly incremental relative risk of 1.09 (CI 1.01 to 1.19, $p = 0.05$). Otherwise stated, this last value translates into an increased risk of perioperative death of 9% greater than the previous year or cumulative risk of 2.48 (CI 1.06 to 5.81, $p = 0.04$) per decade (Table V). Finally, our data

demonstrate that age loses statistical significance when the other risk factors (comorbid state and volume of fluids administered in the first 12 hours postoperatively) are corrected for in a multivariate logistic regression model (OR 2.31, CI 0.84 to 6.38, $p = 0.11$) (Table V). This suggests that age does not act independently of either comorbidity or fluid administration; as the latter is a postoperative event, it may be excluded from a dependent role on age. It stands to reason that older patients are likely to be sicker, and both

factors by themselves but not independently of the other are significant risk factors for perioperative death.

When it was introduced, many investigators hoped to show that preoperative pulmonary function testing would clearly identify those patients at high risk for morbidity and mortality after pulmonary resection. Unfortunately, preoperative values have also lacked consistency as risk factors. Nagasaki, Flehinger and Martini¹⁴ found that a forced expiratory volume in the first second (FEV₁) of less than 40%, a forced expiratory volume in the third second of less than 75%, a diffusing capacity of the lung for carbon monoxide (DLCO) less than 50% and a vital capacity less than 60% of predicted preoperative values were significant risk factors for death. Patel, Townsend and Fountain¹⁰ demonstrated statistical significance only for a ratio of FEV₁/FVC (forced vital capacity) of less than 0.55, whereas others^{5,15} found no value significant in predicting perioperative outcome. Predicted postoperative FEV₁ appears to be more reliable as a risk factor for perioperative death.^{1,17,21,23} as does predicted postoperative DLCO.^{1,18} One study demonstrated a regression model in which the product of predicted postoperative percentage values for FEV₁ and DLCO was the strongest predictor of death after lung resection.¹ Our data included only preoperative values; both measured and percentile of predicted for FEV₁, FVC, FEV₁/FVC, total lung capacity, residual volume and DLCO. None of these were found to exhibit statistical significance between survivor and death groups.

Concomitant medical disease has been identified as a risk factor for perioperative death for many surgical procedures.^{24,25} Following pneumonectomy, comorbidity similarly has been found to be a statistically signifi-

Table III

Death Group Profile

Age, yr	Sex	Side	Cause of death	Autopsy done
73	M	Right	Postpneumonectomy pulmonary edema	No
73	M	Right	ARDS, sepsis	Yes
54	M	Right	Cardiac herniation, diffuse myocardial infarction, 3-vessel disease	Yes
70	M	Left	ARDS, sepsis, respiratory failure, MSOF	No
59	M	Right	Aspiration with secondary ARDS, MSOF, pulmonary embolism	Yes
74	M	Left	Postpneumonectomy pulmonary edema	No
63	F	Left	Pneumonia, acute pericarditis	Yes
64	M	Right	Cardiac herniation	No
80	F	Right	Postpneumonectomy pulmonary edema	No
76	M	Left	Cardiac arrest, ruptured pulmonary artery during cardiopulmonary resuscitation	Yes

ARDS = adult respiratory distress syndrome, MSOF = multisystem organ failure

Table IV

Patient Characteristics and Significant Risk Factors for Perioperative Death After Pneumonectomy Between Survivor and Death Groups

Characteristic/risk factor	Group		<i>p</i> value
	Survivor	Death	
Age, yr*	60.8 (11.2)	68.6 (8.2)	0.04
Sex, M/F	57/25	8/2	NS
Side, left/right	43/39	4/6	NS
Smoking, pack-yr*	49 (40)	44 (23)	NS
Comorbid state*†	0.5 (0.5)	0.9 (0.3)	0.009
Fluids, L (0–12 h)*	1.20 (0.61)	1.61 (0.60)	0.05

*Mean (and standard deviation)

†Reported as a fraction of the study population with 1 or more comorbid states

cant risk factor for death.^{10,11} Our data concur, inferring that sicker patients have a statistically significant increased risk of dying after pneumonectomy such that 50% of survivors and 90% of those who died had 1 or more comorbid states ($p = 0.009$) (Table IV). Univariate logistic regression demonstrated an 8-fold increased relative risk per additional comorbidity (OR 7.92, CI 0.96 to 65.60, $p = 0.05$) (Table V). Multivariate logistic regression analysis showed that, after correcting for age and fluid administration, we were unable to establish statistical significance for concomitant disease (OR 5.27, CI 0.55 to 50.10, $p = 0.15$) (Table V). This suggests that comorbidity cannot be divorced from age; young patients with 1 or more major diseases who undergo pneumonectomy are not at higher risk for death. In other words, senescence may be considered as a comorbid state. Fluid administration may depend on concomitant medical disease states since such patients may be more prone to cardiogenic shock or other states requiring fluid administration.

The volume of intravenous fluids administered perioperatively has been an inconsistent risk factor in the literature, particularly in association with PPE. A significant risk of death correlates with a volume of 3 L or more given in the first 24 hours.¹⁰ Our data (Table IV) demonstrate that the volume of intravenous crystalloid fluid administered in the first 12 hours

postoperatively is a significant risk factor for perioperative death (survivors 1.20 [0.61] L v. death 1.61 [0.60] L, $p = 0.05$). No significant difference was found for the periods of 12 to 24 hours, 24 to 48 hours or 48 to 72 hours after pneumonectomy. Univariate logistic regression demonstrated an OR for death of 2.21/L (CI 0.95 to 5.14, $p = 0.06$) in the first 12 hours postoperatively (Table V). The death rate in our study of patients who received less than 1 L over this period was 2.6% compared with 17.3% in those who received over 1 L. When age and comorbid state were corrected for, the odds ratio for death increased for fluid volume in the first 12 hours while maintaining statistical significance (OR 3.05, CI 1.08 to 8.62, $p = 0.04$), inferring that the volume of fluids administered postoperatively is an important risk factor that functions independently of the other risk factors. Otherwise stated, the young healthy patient who undergoes pneumonectomy has a significant risk of dying should he or she receive excessive fluids in the early postoperative period. The mechanism for this has not been completely elucidated but appears to coincide with our understanding of the pathogenesis of PPE.

First described in cat models by Gibbon and Gibbon in 1942,²⁶ PPE, an almost uniformly fatal complication, is defined as noncardiogenic, noninfectious, non-aspiratory respiratory distress following 2% to 5% of pneu-

monectomies. A rapidly progressive hypoxemia develops with an increased alveolar-arterial gradient and radiographic findings consistent with interstitial and alveolar edema. Although the cause is unclear, the pathophysiology appears multifactorial. After pneumonectomy, the single lung has to accommodate the entire cardiac output and often supernormal flow due to catecholamine release from pain, stress, fever and fluid overload. Increased pulmonary arterial pressure and pulmonary vascular resistance suggestive of pulmonary hypertension have been demonstrated in previous studies.²⁷ Hyperinflation of the remaining lung, which may occur after pneumonectomy, can exacerbate the interstitial accumulation of fluid. In addition, increased capillary permeability has been demonstrated from elevated edema to plasma protein concentration ratios. The lymphatics, already compromised by pneumonectomy, are overwhelmed, especially after right pneumonectomy since fewer lymphatics exist in the left lung. This may be why an increased incidence has been associated with right-sided pneumonectomy. Despite supportive treatment (fluid restriction, diuresis, nutritional support, lower cardiac filling pressures and optimal positive end-expiratory pressure), pulmonary consolidation tends to occur with pneumonia, sepsis and ARDS, and most patients die of multisystem organ failure.

Four retrospective studies have ex-

Table V

Univariate and Multivariate Logistic Regression Analyses

Risk factors	Odds ratio	95% confidence interval	<i>p</i> value	Adjusted odds ratio	Adjusted 95% confidence interval	Adjusted <i>p</i> value
Age, yr	2.48	1.06–5.81	0.04	2.31	0.84–6.38	0.11
Comorbid state*	7.92	0.96–65.60	0.05	5.27	0.55–50.10	0.15
Fluids, L (0–12 h)	2.21	0.95–5.14	0.06	3.05	1.08–8.62	0.04

Multivariate results are corrected for the other 2 risk factors.

*Reported as a fraction of the study population with 1 or more comorbid states

amined the role of perioperative fluid administration in PPE. Zeldin and colleagues reviewed 10 cases and found that the amount of fluids given was the principal factor in developing PPE.²⁸ Verheijen-Breemhaar and colleagues reviewed 243 pneumonectomies over 10 years and noted a 4.5% incidence of PPE, with perioperative fluids constituting a significant risk factor.²⁹ Both studies demonstrated a marked predominance in association with right-side pneumonectomies. Waller and colleagues, in their review of 402 patients, also found no difference in fluids administered.³⁰ Turnage and Lunn reviewed the Mayo Clinic experience of PPE over 11 years and noted that differences between PPE and non-PPE patients with respect to mean perioperative fluids and the volume given in the first 24 hours postoperatively were not significant.²⁷ They demonstrated significant pulmonary hypertension, elevated pulmonary arterial pressures and pulmonary vascular resistance caused by factors other than increased fluid load. Autopsies revealed ARDS as the cause of death in 15 of 17 patients. Mathru and associates measured the edema-to-plasma protein concentration ratio in PPE patients, providing evidence for hyperpermeability of pulmonary capillaries.³¹ This inflammatory response to injury manifested by the lung has previously been identified as the mechanism of ARDS. Although excessive fluid administration may not be necessary for PPE to develop, it is likely contributory in overwhelming the capacity of the remaining lung to remove excess interstitial fluid. Recommendations to prevent the development of PPE include the following: restrict fluids to less than 2.5 L positive balance on the day of operation (including intraoperative fluids), provide adequate analgesia to prevent excessive catecholamine release, limit

airway pressures to avoid overinflation, position the operative side down with the head of the bed elevated to 45° to minimize the gravitational effect on capillary pressure, ensure balanced thoracostomy drainage for 24 to 48 hours to prevent overinflation of the remaining lung as well as mediastinal shift and continue invasive hemodynamic monitoring for 2 to 3 days.²⁷ Prospective studies are needed to determine the efficacy of these recommendations on lowering the incidence of PPE.

CONCLUSIONS

Age, concomitant medical disease and volume of fluids administered in the first 12 hours after pneumonectomy are significant risk factors for perioperative death. Multivariate logistic regression analysis failed to demonstrate independence of age and concomitant disease and fluid challenge remains an independent factor. It is possible that further reduction in the death rate will depend on early postoperative care, specifically in recognizing and preventing PPE.

We appreciate the insightful comments and review provided by Dr. Pierre Ernst, Chairman, Department of Respiratory Medicine, Montreal General Hospital, Montreal.

References

1. Pierce RJ, Copland JM, Sharpe K, Barter CE. Preoperative risk evaluation for lung cancer resection: predicted postoperative product as a predictor of surgical mortality. *Am J Respir Crit Care Med* 1994;150:947-55.
2. Roxburgh JC, Thompson J, Goldstraw P. Hospital mortality and long-term survival after pulmonary resection in the elderly. *Ann Thorac Surg* 1991;51:800-3.
3. Conlan AA, Lukanich JM, Shutz J, Hurwitz SS. Elective pneumonec-

- tomy for benign lung disease: modern-day mortality and morbidity. *J Thorac Cardiovasc Surg* 1995;110:1118-24.
4. Ginsberg RJ, Hill LD, Eagan RT, Thomas P, Mountain CF, Deslauriers J, et al. Modern thirty-day operative mortality for surgical resection in lung cancer. *J Thorac Cardiovasc Surg* 1983;86:654-8.
5. Putnam JB, Lammermeier DE, Colon R, McMurtrey MJ, Ali MK, Roth JA. Predicted pulmonary function and survival after pneumonectomy for primary lung carcinoma. *Ann Thorac Surg* 1990;49:909-15.
6. Harvey JC, Erdman CE, Beattie EJ. Pneumonectomy. *Chest Surg Clin North Am* 1995;5:253-87.
7. Wilkins EW Jr, Scannell JG, Craver JG. Four decades of experience with resections for bronchogenic carcinoma at the Massachusetts General Hospital. *J Thorac Cardiovasc Surg* 1978;76:364-8.
8. Kohman LJ, Meyer JA, Ikins PM, Oates RP. Random versus predictable risks of mortality after thoracotomy for lung cancer. *J Thorac Cardiovasc Surg* 1986;91:551-4.
9. Krowka MJ, Pairolero PC, Trastek VF, Payne WS, Bernatz PE. Cardiac dysrhythmia following pneumonectomy. *Chest* 1987;91:490-5.
10. Patel RL, Townsend ER, Fountain SW. Elective pneumonectomy: factors associated with morbidity and operative mortality. *Ann Thorac Surg* 1992;54:84-8.
11. Romano PS, Mark DH. Patient and hospital characteristics related to in-hospital mortality after lung cancer resection. *Chest* 1992;101:1332-7.
12. Jencks SF, Williams DK, Kay TL. Assessing hospital-associated death rates from discharge data. *JAMA* 1989;260:2240-6.
13. Weiss W. Operative mortality and five-year survival rates in men with bronchogenic carcinoma. *Chest* 1974;66:483-7.
14. Nagasaki F, Flehinger BJ, Martini N. Complications of surgery in the treatment of carcinoma of the lung. *Chest* 1982;1:25-9.
15. Keagy BA, Schorlemmer GR, Murray

- GF, Starek PJK, Wilcox BR. Correlation of preoperative pulmonary function testing with clinical course in patients after pneumonectomy. *Ann Thorac Surg* 1983;36:253-7.
16. Shah R, Sabanathan S, Richardson J, Mearns AJ, Goulden C. Results of surgical treatment of stage I and II lung cancer. *J Cardiovasc Surg (Torino)* 1996;37:169-72.
 17. Wahi R, McMurtrey MJ, DeCaro LF, Mountain CF, Ali MK, Smith TL, et al. Determinants of perioperative morbidity and mortality after pneumonectomy. *Ann Thorac Surg* 1989; 48:33-7.
 18. Ferguson MK, Reeder LB, Mick R. Optimizing selection of patients for lung resection. *J Thorac Cardiovasc Surg* 1995;109:275-81.
 19. Kirsh MM, Rotman H, Argenta L, Bove E, Cimmino V, Tashian J, et al. Carcinoma of the lung: results of treatment over ten years. *Ann Thorac Surg* 1976;21:371-7.
 20. Whittle J, Steinberg EP, Anderson GF, Herbert R. Use of medicare claims to evaluate outcomes in elderly patients undergoing resection for lung cancer. *Chest* 1991;100:729-34.
 21. Bolliger CT, Wyser C, Roser H, Soler M, Perruchoud AE. Lung scanning and exercise testing for the prediction of postoperative performance in lung resection candidates at increased risk for complications. *Chest* 1995;108:341-8.
 22. Breyer RH, Zippe C, Pharr WF, Jen-sik RJ, Kittle CF, Faber LP. Thoracotomy in patients over age seventy years. Ten-year experience. *J Thorac Cardiovasc Surg* 1981;81:187-93.
 23. Markos J, Mullan BP, Hillman DR, Musk AW, Antico VF, Lovegrove FT, et al. Preoperative assessment as a predictor of mortality and morbidity after lung resection [see comment]. *Am Rev Respir Dis* 1989; 139:902-10. Comment in: *Am Rev Respir Dis* 1989;140:1175-6.
 24. Morris JA, MacKenzie EJ, Edelstein SL. The effect of preexisting conditions on mortality in trauma patients. *JAMA* 1990;263:1942-6.
 25. Goldman L, Caldera DL, Nussbaum SR, Southwick FS, Krogstad D, Murray B, et al. Multifactorial index of cardiac risk in noncardiac surgical procedures. *N Engl J Med* 1977;297: 845-50.
 26. Gibbon JH, Gibbon MH. Experimental pulmonary edema following lobectomy and plasma infusions. *Surgery* 1942;12:694-704.
 27. Turnage WS, Lunn JJ. Postpneumonectomy pulmonary edema; a retrospective analysis of associated variables. *Chest* 1993;103:1646-50.
 28. Zeldin RA, Normandin D, Landtwing D, Peters RM. Postpneumonectomy pulmonary edema. *J Thorac Cardiovasc Surg* 1984;87:359-65.
 29. Verheijen-Breemhaar L, Bogaard JM, van den Berg B, Hilvering C. Post-pneumonectomy pulmonary oedema. *Thorax* 1988;43:323-6.
 30. Waller DA, Gebitekin C, Saunders NR, Walker D. Noncardiogenic pulmonary edema complicating lung resection. *Ann Thorac Surg* 1993;55: 140-3.
 31. Mathru M, Blakeman B, Dries DJ, Kleinman B, Kumar P. Permeability pulmonary edema following lung resection. *Chest* 1990;98:1216-8.

**NOTICE OF CHANGE OF ADDRESS
AVIS DE CHANGEMENT D'ADRESSE**

To ensure that you continue to receive the *Canadian Journal of Surgery* without interruption, please fill in and mail this form before you move.

Avant de déménager, assurez-vous de recevoir sans interruption le *Journal canadien de chirurgie* en complétant et en expédiant le formulaire suivant.

Please print / En lettres moulées, svp

Name / Nom

Old address / Ancienne adresse

New address / Nouvelle adresse

Date effective / Date de prise d'effet

Subscribers please mail to: Subscription Administrator, Canadian Medical Association, 1867 Alta Vista Dr., Ottawa ON K1G 3Y6.

Abonnés, veuillez expédier à : Responsable des abonnements, Association médicale canadienne, 1867, prom Alta Vista, Ottawa ON K1G 3Y6.

US address changes / Les Postes américaines enverront les changements d'adresse à : INSA, PO Box 1518, Champlain NY 12919-1518