

Surgical images: soft tissue

Right paraduodenal hernia

A 17-year-old man was admitted to our clinic complaining about intermittent cramping abdominal pain and postprandial nausea of several months' duration. He reported that the pain was worse when supine, especially after the meals, and was relieved in the upright position. He mentioned no previous abdominal operation.

Clinical examination revealed a palpable, slightly tender mass in the epigastrium, which, however, could not be palpated when the patient stood upright. Results from an endo-

scopic examination of the upper gastrointestinal tract were normal. Small-bowel follow-through films taken with diatrizoate meglumine/diatrizoate sodium (Gastrografin®) showed that the jejunum occupied the upper-right part of the abdomen. Multiple jejunal loops were compressed into an ovoid mass situated behind the stomach (Fig. 1). A delay of transit in the jejunum was also documented.

In view of these findings, the patient underwent an exploratory laparotomy, during which a large right paraduodenal hernia was identified

holding almost 1 metre of the jejunum. Afferent and efferent loops of jejunum could be seen entering and leaving the hernia sac. The medial border of the hernia neck consisted of the superior mesenteric and ileocolic artery; the upper and lateral borders were transverse mesocolon (Fig. 2). Exploration of the abdominal cavity revealed no other pathologic findings. The ascending and descending colon were fixated normally on the posterior abdominal wall. The jejunum was easily reduced into the peritoneal cavity and the hernia defect



FIG. 1. Radiograph of the small bowel, showing the position of the jejunal loops behind the stomach.



FIG. 2. Intraoperative view. The hernia neck is clearly visible to the right of the Treitz ligament.

Submitted by Basilio Papaziogas, MD, Anastasio Souparis, MD, John Makris, MD, Athanasios Alexandrakis, MD, and Thomas Papaziogas, MD, PhD, of the 2nd Surgical Clinic, Aristoteles University of Thessaloniki, Thessaloniki, Greece

Correspondence to: Dr. Basilio Papaziogas, Bl. Fanariou str. 16, 551 33 Thessaloniki, Greece; fax 0030 310-992563; papaziog@med.auth.gr

Submissions to *Surgical Images*, soft-tissue section, should be sent to the section editors: Dr. David P. Girvan, Victoria Hospital Corporation, PO Box 5375, Station B, London ON N6A 5A5 or Dr. Nis Schmidt, Department of Surgery, St. Paul's Hospital, 1081 Burrard St., Vancouver BC V6Z 1Y6.

was closed with interrupted sutures.

The postoperative course was uneventful. The patient reported that his preoperative symptoms had not returned.

Right paraduodenal hernias account for about 25% of paraduodenal hernias, the most common type of internal hernia. (Their precise incidence is unknown, since a certain percentage remain asymptomatic throughout the person's lifetime.) These hernias are probably the result of delayed re-entry of the small bowel in the abdominal cavity, leading to its entrapment under the mesentery of the ascending colon.¹ There seems to be a

clear male predisposition, with a ratio of about 3:1.²

Paraduodenal hernias may present with acute intestinal obstruction or intermittent postprandial epigastric pain. Their presence should be suspected without history of previous laparotomy, especially in young patients with intermittent obstructive episodes. Because the herniation of the small bowel through the hernia orifice can be intermittent, preoperative diagnosis is not always possible.^{3,4}

Surgical repair of right paraduodenal hernias includes reduction of the protruding jejunum and closure of the hernia orifice. Large defects of

the mesentery may be covered with use of a mesh.

References

1. Andrews E. Duodenal hernia—a misnomer. *Surg Gynecol Obstet* 1923;37:847-52.
2. Berardi RS. Paraduodenal hernias. *Surg Gynecol Obstet* 1981;152:99-110.
3. Dritsas ER, Ruiz OR, Kennedy GM, Blackford J, Hasl D. Paraduodenal hernia: a report of two cases. *Am Surg* 2001;67:733-5.
4. Yoo HY, Mergelas J, Seibert DG. Paraduodenal hernia: a treatable cause of upper gastrointestinal tract symptoms. *J Clin Gastroenterol* 2000;31:226-9.

Competing interests: None declared.

How you can get involved in the CMA!

The CMA is committed to providing leadership for physicians and promoting the highest standard of health and health care for Canadians. To strengthen the Association and be truly representative of all Canadian physicians, the CMA needs to hear from members interested in serving in elected positions and on appointed committees and advisory groups.

The CMA structure comprises both governing bodies and advisory bodies either elected by General Council or appointed by the CMA Board of Directors. The Board of Directors — elected by General Council — has divisional, affiliate, resident and student representation, is responsible for the overall operation of the CMA and reports to General Council on issues of governance. CMA councils and committees advise the Board of Directors and make recommendations on specific issues of concern to physicians and the public. Four core councils and committees consist of either divisional or regional representation, while other statutory and special committees, and expert working and project advisory groups comprise individuals with interest and expertise in subject-specific fields. Positions on one or more of these committees may become available in the coming year.

For further information on how you can get involved, please contact:

Prunella Hickson
Corporate Affairs
Canadian Medical Association
1867 Alta Vista Drive
Ottawa ON K1G 3Y6
Fax 613 526-7570
Tel 1 800 663-7336, x2249
Email prunella.hickson@cma.ca

By getting involved, you will have an opportunity to make a difference.

We hope to hear from you!